

# POLE BEAN SPECIAL

FOR CONTROL OF: Mexican Bean Beetle, Leafhoppers, Corn Earworms and Nail Head Rust, Uromyces Phaseoli Typica on Pole Beans. Dust once a week. In mountain area when rust is observed or reported, dust twice a week and after every rain using 60 lbs. per acre.

## ACTIVE INGREDIENTS:

Sevin* . . . . .	1.75%
Maneb** . . . . .	3.50%
Sulfur . . . . .	35.00%

INERT INGREDIENTS . . . . . 59.75%

\*1-Naphthyl N-Methylcarbamate.  
\*\*Manganese Ethylenebisdithiocarbamate. 100.00%

**CAUTION:** Make no application on crops later than 4 days before harvest. Harmful if swallowed or inhaled! Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs. Sulfur may burn foliage when temperatures are excessively high.

Do not treat vines that will be fed to dairy animals or animals being finished for slaughter. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop condition, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, handling of this material not in strict accordance with directions given here with.

Reg. U.S. Pat. & T.M. Off.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

**MANUFACTURED BY  
MIKE INC. NEWLAND, N. C.  
DIVISION OF  
SOUTHERN AGRICULTURAL CHEMICALS, INC.  
KINGSTREE, S. C.**