

SEVIN 2% GARDEN DUST

ACTIVE INGREDIENT:
 Carbaryl (1-Naphthyl N-Methylcarbamate)
 INERT INGREDIENTS

2.00%
 98.00%
 100.00%

DIRECTIONS: Beans, Lettuce— For control of MEXICAN BEAN BEETLES, LEAFHOPPERS, FLEA BEETLES— Apply in a manner to thoroughly cover upper and lower leaf surfaces. (In some cases use 10 ounces per 150 feet of row, more or less depending on size of plants). DO NOT TREAT LEAF LETTUCE WITHIN 14 DAYS OF HARVEST. DO NOT TREAT HEAD LETTUCE OR CABBAGE WITHIN 7 DAYS OF HARVEST.

CORN: For control of CORN EARWORMS dust the silks lightly every three days, during silking period. Application of this product to the tassel region of corn during the pollen period will seriously reduce bee population.
 Apply to silks as soon as they appear. Make 3 or 4 applications at 2-3 day intervals as infestation warrants. Apply immediately after rain.

VEGETABLES: For control of Flea Beetles, use at 40-50 lbs. per acre. Treat infested fields before the plants are 10 inches high. When treating during harvest make applications immediately after, rather than before, rain.

POTATOES: For control of LEAFHOPPERS— For control of COLORADO POTATO BEETLES, LEAFHOPPERS, apply in a manner to thoroughly cover upper and lower leaf surfaces.
 For control of JAPANESE BEETLES start dusting when first beetles appear.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

CAUTION: KEEP OUT OF REACH OF CHILDREN

For more information on the use of this product, or to report direct treatment or residues on crops. Protective information may be obtained from the National Agricultural Extension Service.

Precautions: Do not inhale dust. Do not take internally. Avoid breathing dust. Avoid contact with eyes. Wash thoroughly after use. Change contaminated clothing daily. Avoid contamination of food, feed, or other crops that will be fed to dairy animals or animals being finished for slaughter.

EPA REG. NO. 3743-256