



*The Sign of Quality*

POST OFFICE DRAWER 527

**Kingstree, S. C. 29556**

# 10% DITHANE\* DUST

**A Protective Fungicide For The Control Of  
TOBACCO BLUE MOLD**

**ACTIVE INGREDIENTS:**

Zineb (Zinc Ethylene Bisdithiocarbamate) not less than  
(Zinc as metallic 1.5%)

6.5%

**INERT INGREDIENTS:**

93.5%

100.00%

## **CAUTION—KEEP OUT OF REACH OF CHILDREN**

### **DUST TREATMENTS**

**Dust Treatment—Just as Spray Treatments—Should begin before Blue Mold Appears.**

Use a good crank type duster, though small areas can be treated with a small puff-type duster. To make the dust flow evenly, fill the hopper not more than two-thirds full. Apply the dust early in the morning when the air is quite and the plants are moist with dew. The usual plant bed is 5 yards wide or more, and the cotton cover must be removed to dust. Narrow beds 3 or at most 4 yards wide, with side walls that hold the cotton well above the plants can be dusted through this cover. After each treatment all exposed leaf surface should show a coating of dust.

Begin early, with plants the size of a dime, or when blue mold is reported in the area, dust regularly, twice a week, and apply enough dust. The quantities to be applied to each 100 square yards of bed are as follows:

**Applications:**

First to fourth

Fifth to sixth

Seventh and subsequent

Pounds

1 to 1½

2 to 2½

3 to 3½

The 3 to 3½ pound rate is always used during the 2 weeks before transplanting.

When a dust application is washed off by rain while blue mold is active, repeat the treatment at once. If blue mold is seen in a bed that is being dusted, increase the rate and make three instead of two applications that week. Eight to twelve applications will be required.

For the control of Downy Mildew of summer squash and cucumbers, early and late Blight of Tomatoes, dust thoroughly using 30-40 lbs. per acre at weekly intervals and after every rain. Begin applications at first sign of infection. Dust twice weekly under severe blight conditions.

FOR LATE BLIGHT CONTROL on maturing tomato plants use approximately 40 lbs. per acre when applied with ground duster and 50 lbs. per acre when applied with aircraft.

**CAUTION—May cause irritation of nose throat and skin. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Do not reuse container. Destroy when empty.**

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop condition, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA Reg. No. 3743-265