

PHONE 354-6111

Southern

AGRICULTURAL CHEMICALS, INC.



The Sign of Quality

POST OFFICE DRAWER 527

Kingstree, S. C. 29556

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

Application should be made only by experienced personnel using approved safety equipment. Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operations. Do not allow mixture to stand. If mixture does not mix but tends to separate as an oily layer, do not use, as injury to plants may result. Do not combine emulsifiable liquids with wettable powders in the same spray tank, unless previous use of the materials being combined has proved them to be physically compatible. Bury any unused materials at least 18 inches deep.

RECOMMENDATIONS

BEANS: Aphids, Armyworms, Leafhopper, Leaf Miner, Leaf Roller, and Mexican Bean Beetle—Use 1 pt. per acre in sufficient water for thorough coverage. Repeat applications at 7- to 10-day intervals until controlled. Do not apply within 7 days of harvest.

TOBACCO: For control of Southern Potato Weevil and Tobacco Weevil use at rate of 2 qts. per acre. Spray on soil surface and disc in lightly. Treat 1-4 weeks before setting.

COLE CROPS: (Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kale, Mustard, Turnip) Aphids, Armyworm, Diamond-back Moth, Imported Cabbageworm. Treat with 1 pt. per acre in sufficient water for thorough coverage. Do not apply to Kale, Mustard, Turnip within 10 days of harvest. On other crops do not apply within 10 days of harvest.

DESTROY EMPTY CONTAINERS
Perforate, Crush

COTTON: Aphids, Red Spider Mites—Use 1/2 pt. per acre in sufficient water for thorough coverage. Do not apply within 7 days of harvest.



CUCURBITS: (Cucumber, Squash and Melons): Aphids, Cucumber Beetle, Melonworm, Pickleworm, Serpentine Leaf-miner—Use 1 pt. per acre in sufficient water for thorough coverage. Do not apply unless plants are dry nor until runners begin to form. Do not apply within 15 days of harvest on cucumbers and squash; 7 days on melons.

POTATO, TOMATO: Aphids, Flea Beetle, Leafhopper, Serpentine Leaf Miner—Use 1/2 pt. per acre in sufficient water for thorough coverage. Do not apply within 10 days of harvest on tomatoes and 5 days of harvest on potatoes.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop condition, mixtures with other chemicals not specifically recommended, and other interfering factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

This product will kill fish and wildlife. Keep out of any body of water. Apply only as specified on this label.

READ ALL DIRECTIONS CAREFULLY BEFORE USING. For more information, see back panel.



DANGER—POISON—PRECAUTIONS



POISONOUS IF SWALLOWED

This product can kill you if swallowed even in small amounts. Spray mist or dust may be fatal if swallowed.

POISONOUS BY SKIN CONTACT

Poisonous if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes).

POISONOUS IF BREATHED

Breathing vapors, spray mist or dust may be fatal.

POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur.

Consult your State Agricultural Extension Service or Experiment Station regarding treating treated areas.
CONTAINER DISPOSAL: See back panel for proper disposal.
CROP USE DIRECTIONS: Use only according to back panel instructions.

REST

For retail sale under their di Certified App

STOP

POIS

See side 1 for anti and preca

PA

ACTIVE Parat Aron. INERT IN

NET VOLU

South

NOTE TO PHYSICIAN:
Antidote for water-alcohol and late in anaphylaxis. TWO TO FOUR mg intravenously or intramuscularly as soon as signs of shock occur. Repeat at 10 to 15 minute intervals until symptoms of anaphylaxis disappear. 2 PMACs in 24 hours are safe and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Paralysis is a static state; therefore, that for affecting the central and peripheral nervous systems and producing, and an and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued administration of the drug may cause severe and fatal relapses have been reported. The drug should be given under close supervision of the patient indicated for at least 48 hours.

EPA Reg. No. 3743 260
E.P.A. EST. 3743 SC-1