


10% Sevin* Dust



ACTIVE INGREDIENT

Carbaryl (1-Naphthyl Methylcarbamate) _____

10.00%

INERT INGREDIENTS _____

90.00%

TOTAL _____

100.00%

*Trademark of Union Carbide Chemical Co. for 1-Naphthyl N-Methylcarbamate

CAUTION: Keep Out of Reach of Children

soon as plants are set and repeat as needed. Dust thoroughly particularly undersides of leaves for flea beetle control.

Cotton: To control Boll Weevil, Bollworm, Cotton Leafworm, Lygus Bugs, Cotton Leaf Perforator, Thrips, and Cotton Fleahoppers follow State recommendations for frequency and timing of applications. Usually for early season control, two to three applications should be made at 7 to 10 day intervals at the rate of 10 pounds per acre. For mid-season control, apply 15 pounds per acre. Repeat application at 5 to 7 day intervals until infestation is brought under control. Usually four applications will be necessary. Repeat a series of four applications at 5 to 7 day intervals if boll weevils reappear in sufficient numbers to cause damage. For late season control, apply at the rate of 20 pounds per acre at 5 to 7 day intervals. Anytime bollworms are a threat, when as many as 4 to 5 small worms and eggs are found per 100 terminals examined, apply at the rate 15-20 pounds of dust per acre. Direct the dust down ward into the tops of the plant where the worms are hatching. Repeat applications at 5 day intervals until infestation is controlled. The early season dosages will control only thrips, cotton fleahoppers, and cotton leafworms. Suppression of aphid populations may be obtained by repeated applications.

CAUTION: HARMFUL IF SWALLOWED OR INHALED! Do not take internally. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and food stuffs. Do not apply after bolls open. Do not graze treated fields or feed trash to dairy or meat animals. Wear regular long sleeved work clothing. Wash hands and face before eating.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative agricultural extension service.

DO NOT REUSE EMPTY CONTAINER. DESTROY BY BURYING IN A SAFE PLACE.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop condition, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

E.P.A. Reg. No. 3743-277

Manufacturers name, address and net contents printed on container.