



NC.

OFFICE DRAWER 527

tree, S. C. 29556

# IODAN MISCIBLE

## INSECTICIDE

### ACTIVE INGREDIENTS

Iodan (Hexachlorohexahydromethano-2, 4, 3- benzodioxathiepin oxide)

24.00%

69.00%

7.00%

100.00%



### OTHER INGREDIENTS

U. S. Pat. No. 2,750,005

Iodan Miscible contains 2 parts of Thiodan per gallon.

**WARNING:** Iodan is a contact poison and is absorbed through the skin. Do not breathe spray mist. Do not use in enclosed areas. In case of contact, immediately remove contaminated clothing and wash thoroughly with soap and water. If swallowed, get medical attention. Wash thoroughly with soap and water. Do not use on or near clothing. During commercial or prolonged exposure, use appropriate protective equipment, including synthetic rubber gloves and a mask or respirator of a type approved by the U. S. Dept. of Mines or the U. S. Dept. of Health. Do not apply or allow to drift to areas occupied by beneficial animals. This product is poisonous to fish and wildlife. Keep out of any body of water. Use this product only as specified on this label. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not use on or near water. Clean equipment or disposal of wastes. This product is toxic to fish and wildlife. Do not use on or near water. Workers entering treated area within 24 hours should wear protective clothing.

### WARNING: KEEP OUT OF REACH OF CHILDREN

Destroy by burning in a safe place"

### WARNING—FLAMMABLE

Keep container closed. — Leaking packages should be opened in a well-ventilated area.

### DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY

USE CAREFULLY. DO NOT USE THE MATERIAL IN A MANNER OR AT A TIME OTHER THAN THAT SPECIFIED ON THE LABEL. OVERUSE OR MISUSE MAY PRODUCE PLANT INJURY. EXCESSIVE RESIDUES MAY BE DETECTED BY PLANT TESTS.

Manufactured by

Manufactured By

Southern Agricultural Chemicals, Inc.

KINGSTREE, S. C.

**COTTON:** Boll Weevil—1 to 2½ qts. per acre; Bollworm, Cabbage Looper, Cotton Stink Bug, Lygus Bugs, Cotton Leaf Perforator—2 qts. per acre. Do not apply after buds open. Do not graze meat or dairy animals in treated fields.

**CUCUMBERS (C):** Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle—1 to 2 qts. per acre. May be applied up to day of harvest.

**EGGPLANTS (E):** Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Stink Bug, Green Stink Bug, Whitefly—2 qts. per acre. Do not apply within 14 days of harvest.

**GRAPES (G):** Grape Leafhoppers—1 qt. per 100 gallons, 2 to 2½ qts. per acre. Do not apply to Concord variety of grapes as severe injury is likely to occur.

**POTATOES (P):** Aphids, Colorado Potato Beetle, Plant Bugs (including Shield Shaped Bug), Potato Flea Beetle, Potato Leafhopper, Potato Tuberworm, Southern Armyworm—1 to 2 qts. per acre; European Corn Borer, Potato Psyllid—1½ to 2 qts. per acre; Late Chalk Bug—2 qts. per acre; Three-lined Potato Beetle—1 qt. per acre.

**SQUASH, SUMMER AND WINTER (S):** Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle, Squash Vine Borer—1 to 2 qts. per acre. May be applied up to day of harvest. For vine borer control, apply weekly to flower buds. Do not apply when moths first appear.

**NOTICE:** Follow directions carefully. Timing and method of application, whether or not mixed with other chemicals or specifically recommended, and other precautions in the use of this product are beyond the control of the manufacturer. Assume all risks of use, storage or handling of this material, not in any way, in any direction given herewith.

NET WT. 5.0 LBS.