

DIRECTIONS FOR USE

APPLES: For the control of Scab, Powdery Mildew, Codling Moth, Plum Curculio, Orange Tortrix, San Jose Scale, Forbes Scale, Scurfy Scale, Red Banded Leaf Roller, Apple Maggot, Woolly Apple Aphid, Bud Moth, Fruit Tree Leaf Roller, Rosy Apple Aphid, Green Apple Aphid, Leafhopper, European Red Mite and Two-spotted Mite. Apply 50-65 lbs. per acre at 7 day intervals or as needed to maintain control from pre-bloom period through first cover. Repeat at 7-10 day intervals throughout the summer months. Make full coverage applications. Control of scale insects is limited to the crawler stage only. Do not apply within 14 days of harvest. Do not apply to susceptible varieties such as McIntosh, Cortland, and related varieties. Golden Delicious, Grimes Golden, Rhode Island Greening or Jonathan. Do not use treated apples in the manufacture of apple products for use in livestock feeds.

USE PRECAUTIONS

Do not prune fruit trees or thin fruit within 15 days after application. Do not apply during bloom when honey bees or other pollinators are active. Avoid contact with dust drift on flowers, shrubs, or fruits, vegetables and forage crops. Keep children and animals off treated areas until chemical has been washed into the soil by sprinkling or rain and the grass is dry. Do not graze cover crops in treated orchards.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and do not apply closer to harvest than specified. Consult your State Agricultural Experiment Station for specific information consistent with labeling as the timing, number and rate of applications may vary with local conditions.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.



DANGER—POISON—PRECAUTIONS

POISONOUS IF SWALLOWED

This product can kill you if swallowed even in small amounts: spray mist or dust may be fatal if swallowed.

POISONOUS BY SKIN CONTACT

Poisonous if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes.)

POISONOUS IF BREATHED

Breathing vapors, spray mist or dust may be fatal.

POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

POST TREATED AREA

Consult your State Agricultural Extension Service or Experiment Station regarding posting treated areas.

CONTAINER DISPOSAL

Burn empty bags, stay out of smoke. This product is to be used by the commercial growers or commercial applicators.

STOP - READ LABEL BEFORE USE

POISON

See side panels
for antidote
and precautions



CAN KILL YOU

CYPREX, PARATHION, SULPHUR

2-2-20 DUST

NOT FOR HOME USE

ACTIVE INGREDIENTS:

Dodine (n-Dodecylguanidine acetate)
Parathion O, O-Diethyl O-p-Nitrophenyl
Thiophosphate)

Sulphur
INERT INGREDIENTS.

NET WEIGHT


MANUFACTURED BY

Southern Agricultural Chemicals, Inc.

KINGSTREE, S. C.

NEWLAND, N. C.

Cyprex is a registered trademark of American Cyanamid
U. S. Patent No. 2,867,562



3743-283

