

DIPEL® 150 DUST*

(Bacillus thuringiensis)

ACTIVE INGREDIENT:

Bacillus thuringiensis, Berliner, Potency of 320 International Units per mg. (at least 0.5 billion viable spores per gm.)*

0.064%

INERT INGREDIENTS

99.936%

Total

100.00%

*Equivalent to potency of 0.15 billion International Units per pound of this product.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

REASONABLE CARE SHOULD BE EXERCISED TO AVOID INHALATION OR CONTACT WITH OPEN WOUNDS. DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY.

DIRECTIONS FOR USE

Dipel 150 Dust is a Biological Insecticide intended for use against Chewing Larvae (Caterpillars) of Certain Lepidopterous Species.

Dipel 150 Dust is recommended for use by ground or by Air.

Apply when pests are first noticed and repeat as necessary.

Thorough Plant Coverage is necessary for control.

APPLICATION RATE

For the control of Cabbage Looper and Imported Cabbage Worm on Broccoli, Cabbage, Cauliflower, Collards, Kale, Mustard Greens, Turnip Greens. Apply 25 to 40 lbs. per acre.

For the control of Cabbage Looper on Celery, Lettuce, Melons, Potatoes, Spinach, Tomatoes and Cucumbers (Southeastern U. S. Only). Apply 25 to 40 lbs. per acre.

For the control of Hornworm on Tobacco and Tomatoes. Apply 12 to 25 lbs. per acre.

®Dipel is the registered trademark of Abbott Laboratories, North Chicago, Illinois.

MANUFACTURED BY

SOUTHERN AGRICULTURAL CHEMICALS

KINGSTREE, S. C.

NEWLAND, N. C.

NET CONTENTS

E.P.A. Reg. No. 3743-289

JUN 1974
UNDER THE FEDERAL
INSECTICIDE ACT
7/1/74