

MANEB DUST ³⁷⁴³⁻²¹⁴

ACTIVE INGREDIENTS:

Maneb (Manganese ethylenebis(lithiocarbonate))

6.0%

INERT INGREDIENTS;

94.0%

³⁷⁴³⁻⁴⁴
100.0%

Caution - Keep Out Of Reach Of Children

DIRECTIONS

Use at the rate of 20 to 30 pounds per acre, dusting thoroughly to give coverage to undersides of leaves.

TOMATOES — For control of Early Blight and Late Blight, Anthracnose, Septoria Leaf Spot and Gray Leaf Spot, start application when first fruit clusters are formed, repeat at 7 to 10 day intervals. Do not apply within 5 days of harvest.

POTATOES — For control of Early Blight and Late Blight start application as soon as plants are 2 to 6 inches high. Repeat at 5 to 10 day intervals as long as blight threatens.

CUCUMBERS AND SQUASH — For control of downy mildew and anthracnose, start treatments as soon as plants begin to run or disease threatens and repeat every 5 to 10 days as long as needed. Do not apply within 5 days of harvest.

SPINACH — Downy mildew (Blue Mold) start applications as soon as disease appears in area and continue at 7 to 10 day intervals. Do not apply within 10 days of harvest.

CAUTION — Avoid breathing of dust. Avoid contact with skin. Wash thoroughly. Avoid contamination of feed and foodstuffs.

May cause irritation of eyes, nose, throat and skin. May be harmful if inhaled or swallowed. In case of contact, flush with plenty of water. For eyes, get medical attention immediately. Remove harmful pesticidal residues at harvest from edible parts of growing food crops by washing, brushing or other means. Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

E.P.A. Reg. No. 3743 - 296