




POST OFFICE DRAWER 527
Kingstree, S. C. 29556

3743-298
PM12
171

BEST AVAILABLE

BEST AVAILABLE COPY

Contains 4-lb. of Methyl Parathion per gallon and is an emulsifiable concentrate that mixes readily with water.



DIRECTIONS FOR USE

Crop: Cotton
Diseases: None
Insects: Aphids, Boll Weevil, Cotton Leaf Perforator, Leaf Worms, Spindles, Mites

Immediately prior to use mix the material with water. A good practice is to dilute the material with just to exceed an equal amount of water, and then add the diluted emulsion to the required amount of water. Agitate the material continuously while mixing and applying.

Application of this material should be made when an infestation reaches the point where control is required. Full coverage of plants during applications is essential to good control. Use Methyl Parathion 4 Emulsifiable at the rate of 2 pints per acre in sufficient water to provide uniform coverage. Application should be made at 5 to 7 day intervals depending on the severity of the infestation and local conditions.

Methyl parathion will control the *Trialeurodes* and *Desmetorium* species of spider mites but will not control the *Bimaculatus* (two spotted) species.

Workers entering treated fields within 24 hours after application should wear protective clothing.

THOROUGH COVERAGE IS NECESSARY

Do not apply on cotton within 15 days of hand picking.
Cotton which has received late applications of this insecticide should not be grazed by dairy animals or animals being finished for slaughter.

For specific uses or for local conditions, it is suggested that helpful information may be obtained from your State or Federal Agricultural Authorities who have had experience with this type of material.



DANGER—POISON—PRECAUTIONS

POISONOUS IF SWALLOWED



POISONOUS BY SKIN CONTACT



POISONOUS IF BREATHED

Breathing vapors, mist or dust may be fatal.

KEEP OUT OF REACH OF CHILDREN AND WILDLIFE
Do not use in areas where children, birds, bees, and other animals may be present. Do not use in areas where crops are being raised. Keep out of reach of children. Do not use in areas where crops are being raised. Do not use in areas where crops are being raised. Do not use in areas where crops are being raised.

Do not use in areas where crops are being raised.

RESTRICTED USE PESTICIDE
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

STOP - READ LABEL BEFORE USING!

POISON
See side panels for antidote and precautions



DANGER
Keep out of reach of children

CAN KILL YOU

METHYL PARATHION 4 EMULSIFIABLE

NOT FOR HOME USE

ACTIVE INGREDIENTS

Methyl Parathion 4 Emulsifiable	15.00%
Petroleum hydrocarbons	47.00%
Inert ingredients	8.00%
Methyl Parathion	100.00%

Manufactured By
Southern Agricultural Chemicals, Inc.
KINGSTREE, S. C.

USE ONLY WHEN WEARING THE FOLLOWING:

1. Long-sleeved shirt and long trousers or pants.

2. Closed-toe shoes.

3. Goggles or safety glasses.

IMPORTANT: Do not use if you are pregnant or nursing. Do not use if you are allergic to any of the ingredients. Do not use if you are taking any medicine.

POISON SIGNS

Reddening of the face, nausea, vomiting, diarrhea, dizziness, headache, and weakness.

FIRST AID

Call a doctor, physician, clinic or hospital if exposed to parathion and describe his condition. If breathing has stopped, start artificial respiration. If swallowed and victim is awake, encourage the victim to drink down throat or by giving the victim a glass of water. Never give anything by mouth to an unconscious person. See doctor immediately. In case of contact, immediately flush eyes or skin with running water, removing contaminated clothing and shoes.

NOTE TO

Antidote administered: atropine sulfate in 1 mg. increments as soon as symptoms of overdose are observed. Repeat 2 PAM. Provide a glass of water with atropine. DO NOT GIVE M. P. H. U. if necessary, as it will affect the central nervous system and respiratory depression. At least 6 cc supplemental oxygen and treated symptoms may occur and fatal reactions have been reported. SUPERVISION OF THE PATIENT IS INDICATED.

NOTICE: Follow directions carefully. Timing conditions, mixtures with other chemicals, and other factors in the use of this product are beyond the scope of this label. For full details of use, storage, and handling of this material, see the product literature.