

# ROYAL BRAND 20% LINDANE EC

CONTAINS 1.6 LBS. TEC. LINDANE PER GALLON

|   |      |
|---|------|
| ACTIVE INGREDIENTS                                  |      |
| Gamma Isomer of benzene hexachloride (from Lindane) | 20%  |
| Petroleum Hydrocarbons                              | 74%  |
| INERT INGREDIENTS                                   | 6%   |
| TOTAL   | 100% |

## CAUTION: KEEP OUT OF REACH OF CHILDREN

For squash vine borer, striped and spotted cucumber beetles, melonworms, pickleworms and aphids on cucurbits use at the rate of 2 pint 20% Lindane per acre. Begin application on foliage when worms first appear and repeat weekly.

For Balsam woolly aphid use 2 quarts of 20% Lindane EC per 100 gallons of diesel oil and spray base of tree 18 inches up, higher if there are attacks. Spray entire tree during May, June, or July. IN CASE OF DANGER TO SHRUBBERY USE WATER SUBSTITUTED FOR OIL.

For Turpentine beetle mix 12 tablespoons of 20 % Lindane EC with one gallon of diesel oil. Keep trees vigorous and free from injury to trunk. Avoid excessive grading or filling. Cut dead trees. Prevent or control attack by spraying lower 12 feet of trunk. Spray injured trees as precaution. Do not apply on wet bark. USE SPRAYS DILUTED WITH WATER ONLY NEAR SHRUBBERY OR LAWNS.

CAUTION: Do not use Lindane in fields to be planted later in potatoes or other root crops, as it may adversely affect their flavor.

CAUTION: May be absorbed through skin. Avoid inhalation and skin contact. In case of contact, wash immediately with soap and water. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Harmful if swallowed. Keep out of reach of children. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.


This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop condition, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Manufactured By

**SOUTHERN AGRICULTURAL CHEMICALS, Inc.**  
KINGSTREE, SOUTH CAROLINA

E. P. A. Reg. No. 3743-302  
E.P.A. EST. NO. 3743-SC-1



29556

TR 527

PHONE 354-6111

*Southern*  
AGRICULTURAL CHEMICALS, INC.