

GENERAL DIRECTIONS

1. Add about half the total amount of water to be used to the spray tank. Start motor or pump bypass and add proper amount of COTTON-TOX MP-6-15. Continue mixing and add remaining water.

DIRECTIONS FOR USE ON COTTON:

COTTON-TOX MP-6-15 may be applied by either ground equipment or aircraft. For ground equipment, dilute in sufficient water to give adequate coverage. For aerial applications, dilute in 3 to 4 gallons of water per acre.

EARLY SEASON PROGRAM: For general early season control, including plant bugs and fleahoppers, use 1.5 pints COTTON-TOX MP-6-15 per acre. Make two or three applications at 7 day intervals starting at the two-leaf stage. For thrip control only, use 0.75 to 1 pint per acre and for cutworms use 3 pints per acre.

MID SEASON INSECTS: Overwintering boll weevils, plant bugs, aphids and cutworms are controlled with 2 to 3 pints per acre of COTTON-TOX MP-6-15. Repeat treatments at 7 day intervals as insect counts indicate. Repeat as necessary.

LATE SEASON CONTROL: Use 3 quarts per acre to control plant bugs, bollworms and aphids. Use the higher rate for heavy bollworm outbreaks. Repeat applications at 5 to 7 day intervals as insect counts indicate. Repeat as necessary.


CAUTION: Do not backpack or harvest cotton in treated areas. Do not allow children within 48 hours of treatment. Do not wear treated clothing. Do not permit children to play in treated areas. Do not allow for slaughter in treated field.


DIRECTIONS FOR USE ON SOYBEANS:

COTTON-TOX MP-6-15 controls plant bugs, cutworms, and aphids, and is effective against corn earworms, soybean leafhoppers, and soybean cutworms. Apply 1.5 pints per acre.

For control of plant bugs, cutworms, and aphids, apply 1.5 pints per acre. For control of corn earworms, soybean leafhoppers, and soybean cutworms, apply 3 quarts per acre.

ACCEPTED
AUG 3 1973
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR CHEMICALS LISTED REGISTERED
 UNDER THE FEDERAL INSECTICIDE
 ACT



Dangerous Poison
 Freely Available


POISONOUS IF SWALLOWED

This product can kill you if swallowed even in small amounts; spray mist or dust may be fatal if swallowed.

POISONOUS BY SKIN CONTACT

Poisonous if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes.)

POISONOUS IF BREATHED

Beathing vapors, spray mist or dust may be fatal.

POISONOUS TO FISH & WILDLIFE

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

POST TREATED AREA

Do not enter treated areas until advised by the Agricultural Extension Service.

CONTAINED IN

1.5 Pint Containers
 3 Quart Containers
 5 Gallon Containers

STOP - READ

POISON
 See side panels
 for antidote
 and precautions

CA

COTTON

ACTIVE INGREDIENTS
 •Toxaphene
 •O, O-dimethyl Xylene (Aroma)
INERT INGREDIENTS

•Technical chlorinated Methyl Parathion

Used as directed by the manufacturer against bollworms, Plant Bugs, Fleahoppers, Cotton Leafworms, Green Cutworms, and Cotton Leaf Caterpillars.

See antidote information on side panels. DO NOT HEAT OR OVER FILL.

Southern Agricultural Chemicals

1000 West 10th Street
 Fort Worth, Texas 76102

Manufacturer's name, address and net contents printed on container.

