

3743-331

ROYAL BRAND
Dairy and Livestock Dust
FOR HORNFLIES, LICE AND FACE FLIES

**CAUTION: KEEP OUT OF REACH
OF CHILDREN**

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:

Methoxychlor, Technical* 5%
Malathion (O, O-dimethyl dithiophosphate of
diethyl mercaptosuccinate) 4%

INERT INGREDIENTS:

91%
100%

*Equivalent to 4.1% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 0.6% related com-
pounds.

MANUFACTURED BY

**SOUTHERN AGRICULTURAL
CHEMICALS, INC.**

Kingstree, S. C.

Newland, N. C.

DIRECTIONS FOR USE

FOR DIRECT APPLICATION TO DAIRY CATTLE AND BEEF CATTLE, FOR CONTROL OF HORNFLEES:

Apply to the horns and base of horns of cattle. Rub lightly into hair. Repeat application in 10 to 14 days if necessary. Do not repeat more often than once in every three week period.

FOR CONTROL OF LICE AND TICKS ON BEEF CATTLE

Treat entire animal very thoroughly. Repeat treatment if necessary. May be used on non-lactating dairy animals if necessary with care when freshening.

AS AN AID FOR THE CONTROL OF FACE FLIES ON BEEF CATTLE

Put the faces of cattle through spray. Repeat as necessary.

FOR CONTROL OF HOG LICE

Apply to hogs, pens, and pits. Repeat treatment as necessary. Repeat in 10 days or there after as needed.

LICE AND FLIES ON SHEEP

Apply to animals thoroughly for control of lice and flies. Repeat as necessary.

CAUTION

Do not use on horses, dogs, cats, birds, or other animals. Do not apply to the face of humans. Avoid contact with eyes. Wash thoroughly after use.

Keep out of reach of children. Do not use on animals with open wounds.

NOTE: For information on the use of this product, consult the label. Do not use on animals with open wounds. For more information, contact your local health department.

1974-1975