

1 OTH E CHAMER 127

Kingstree, S. C. 29556

ROYAL BRAND TOMATO DUST

ACTIVE INGREDIENT

Copper (in basic copper sulphate expressed as metallic)
INERT INGREDIENTS

7.00% **93.00%** 100.00%

A FUNGICIDE DUST FOR USE ON

TOMATOES and POTATOES for the control of early and late blights and as an aid in controlling bacterial spot. Dust plants thoroughly at weekly intervals and following rains. For potatoes make 30 to 50 lbs. per acre applications beginning when plants are 6 inches high. For tomatoes make 30 to 50 lbs. per acre applications beginning when plants are set in the field.

CUCUMBERS, SQUASH, WATERMELONS and CANTALOUPES for the control of downsy mildew. Dust thoroughly, but lightly, using 15 to 40 lbs. per acre. Apply at 10-lay intervals and following rains.. Heavy applications, applications to tender

foliage, and applications to wet foliage can cause injury to cucurbits.

Make first application as soon as leafs appear or as soon as vines begin to run.

PEITER as an aid in the control of bacterial spot, apply at weekly intervals and if possible preceding heavy rains, using 20 to 45 lbs. per acre applications. First application when spots appear.

There is no time limitation on the use of this dust. Remove excess residue at harvest. Consult proper State Extension specialist to determine if local recommendations are consistent with this label before using. Do not exceed indicated dosages. May cause injury to tender plant parts particularly when applications are excessive, uneven, or when foliage is wet. Control of the indicated diseases may not be satisfactory during periods of frequent or heavy rainfall, when disease infestations are extremely severe, or when applications are uneven, inadequate, or poorly timed.

Do not apply or allow drift to corn, grain sorphum, or other copper sensitive plants. Do not apply when conditions favor drift from treated area.

Do not reuse empty container. Destroy it by burying with west or burning. Stay away from smoke or funes.

KEEP OUT OF REACH OF CHILDREN

CAUTION Harmful if swallowed. Avoid contact with skin, eyes, and clothing. Do not contaminate feed or foodstuffs.

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with the directions given herewith.

MANUFACTURED BY

Southern Agricultural Chemicals,

KINGSTREE, S. C.

NEWLAND, N. C.

E.P.A. EST NO. 3743-8C-1 E.P.A. REG. NO. 3743-334

