


Southern AGRICULTURAL CHEMICALS, INC.

POST OFFICE DRAWER 527
Kingstree, S. C. 29556



APR 1975
U.S. Dept. of Agriculture
EPA Reg. No. 3743-336

DIRECTIONS

M-M 2-4 Cabbage Dust is recommended for control of certain insects on cabbage and downy mildew. Apply with standard dust application equipment. Use only in commercial plantings; do not use in home plantings.

Application: Apply at the recommended rates when insects and downy mildew first appear.

Continue applications at 5- to 7-day intervals or as needed. Apply the low rates on small plants, small worms and light infestations of insects. Use intermediate rates on larger worms and heavier infestations of insects. Use 1 to 3 applications of the highest recommended rate for controlling severe infestations. Thereafter, use the lowest rate possible to maintain control. The number of days in parentheses following the dosage rates is the preharvest interval which must be allowed between the last application and harvest. For downy mildew begin applications 7 - 10 days after planting with repeat applications at 3 - 5 day intervals.

Cabbage — Cabbage Looper, diamondback moth and downy mildew 30 - 45 lbs. / per acre (7 day).

IMPORTANT

Not for use or storage in or around the home. Keep animals and unprotected persons out of operational areas during treatment and while there is danger of drift.

Do not re-use container; destroy when empty. Do not contaminate water when cleaning equipment or disposing of wastes or containers.

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply when runoff is likely to occur.

Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

ROYAL BRAND M-M 2-4 CABBAGE DUST

For Control of Certain Cabbage Insects.
and DOWNEY MILDEW

ACTIVE INGREDIENTS: Methomyl S-methyl-N- ((methylcarbamoyl) oxy) thioacetimidate.....	2%
Maneb (manganese ethylenebisdithiocarbamate)	4%
INERT INGREDIENTS	94%

WARNING: KEEP OUT OF REACH OF CHILDREN WARNING! HARMFUL OR FATAL IF SWALLOWED OR INHALED

- Do not breathe dust.
- Do not get in eyes, on skin, on clothing.

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention; remove and wash contaminated clothing before re-use.

WEAR PROTECTIVE CLOTHING
Wear goggles and a mask or respirator of a type suitable for protection against insecticide dust when loading, applying or otherwise handling. Wear clean clothes daily. Wash thoroughly after handling and before eating or smoking. Do not contaminate or store near feed or foodstuffs.

ANTIDOTE AND FIRST AID

ATROPINE IS AN ANTIDOTE-CONSULT PHYSICIAN FOR EMERGENCY SUPPLY OF 1/100 GRAIN ATROPINE TABLETS. CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

If swallowed, give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. If inhaled, remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately. **Never give anything by mouth to an unconscious person.**

If warning symptoms appear (See warning symptoms below), before physician arrives, immediately swallow two atropine tablets (each 1/100 gr.); thereafter, every 10 to 15 minutes, take one atropine tablet (1/100 gr.) until throat becomes dry and skin becomes dry and flushed. Take additional tablets as necessary to maintain a moderately dry throat and dry, flushed skin until physician arrives.

NOTE TO PHYSICIAN

Warning Symptoms — Methomyl poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in the chest, constriction of pupils, sweating, slow pulse, muscle tremors.

Treatment — Atropine sulfate should be used for treatment. Administer repeated doses, 1.2 to 2.0 mg. intravenously every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until the patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured.

Do not use morphine or 2-PAM.

NOTICE: Seller makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material except those which are contained on Seller's label.

Apply This Product Only as Specified on this Label.

E. P. A. Reg. No. 3743-336
E.P.A. EST. NO. 3743-SC-1