

CUKE SPECIAL

ACTIVE INGREDIENTS:

Maneb (Manganese Ethylenedisithiocarbamate) ...
Gamma Isomer of Benzene Hexachloride (from Lindane) ...

INERT INGREDIENTS

ACCEPTED

APR 21 1975

Under the Federal Insecticide,
Fungicide, and Plant Repellent Act
as amended, for use as pesticide
registered under

DIRECTIONS: No. 3743-339

48%
10%
42%
100%

For control of pickleworms, downy mildew and anthracnose on cucumbers. Use at rate of 6 1/4 lbs. per acre. Spray 7-10 day intervals with sufficient water to give thorough coverage. Begin when diseases threaten or when plants begin to run.

DO NOT APPLY WITHIN 5 DAYS OF HARVEST

CAUTION - KEEP OUT OF REACH OF CHILDREN

CAUTION. Avoid contact with skin and eyes. Avoid breathing dust. Do not contaminate feed or foodstuffs. Keep away from children and domestic animals. Wash after handling, may be absorbed through skin.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply when runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residue on crop. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop condition, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material in any event in accordance with directions given herewith.

E. P. A. REG. NO. 3743-339

E. P. A. ISI NO 3743-SC-1

29556

PHONE 394.6111

Manufacturers name, address and not contained in label or container.

Southern