

GENERAL DIRECTIONS

CIODRIN® Insecticide Emulsible Concentrate is suitable for use in conventional hydraulic sprayers or low pressure knapsack sprayers. Just prior to use, mix the required amount of CIODRIN® Insecticide Emulsible Concentrate in sufficient water and apply in accordance with State and Federal Recommendations. For control of stable flies, horn flies, face flies, house flies, lice and louse star ticks on dairy cattle, beef cattle, sheep, goats and hogs.

DIRECTIONS FOR LIVESTOCK PEST CONTROL

Mix 1 quart of CIODRIN® Emulsible Concentrate (2 lb/gal) per 6 gallons of water (1% concentration). Spray to thoroughly cover all parts of the animal, including the legs, using 1-2 pints of spray per large animal and proportionately less for smaller animals.

For pen or corral operations involving spray applications to livestock with high pressure equipment, do not exceed a 0.5% concentration. Mix 1 quart of CIODRIN® Emulsible Concentrate (2 lb/gal) per 12 gallons of water (0.5% concentration). Spray to thoroughly cover the animal's body and legs (avoid excessive spraying of the head) using 1-2 quarts of spray per large animal and proportionately less for smaller animals.

Improved face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gallons of spray. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

Low volume application for control of face fly

Mix 2 tablespoons (1 fluid ounce) of CIODRIN® Emulsible Concentrate (2 lb/gal) per 1½ pints of water (1% concentration). Using calibrated sprayer, apply 5 ml. of spray to the face and a total of 5-10 ml. to the back and sides of the animal 3 times per week.

For control of house fly, face fly and stable fly in and around livestock buildings.

Mix 1 quart of CIODRIN® Emulsible Concentrate (2 lb/gal) per 6 gallons of water (1% concentration). Spray premises such as inside and outside barn walls (including dairy barns), eaves, fences and other resting or breeding places at least twice monthly using approximately 1 gallon of finished spray per 1000 sq. ft.

BACK-RUBBER APPLICATIONS

For use in back-rubbers or face-rubbers, for the control of horn flies and face flies, mix 1 quart of CIODRIN® Emulsible Concentrate (2 lb/gal) per 7 gallons of back-rubber base oil (1% concentration). This may be diesel fuel or combinations of diesel fuel and motor oil in accordance with local recommendations.



CIODRIN® INSECTICIDE

Emulsifiable Concentrate

CAUTION: (Contains 2.1 lb. Ciodrin Insecticide Per Gallon) KEEP OUT OF REACH OF CHILDREN.
WARNING: SEE SIDE PANEL FOR ADDITIONAL INFORMATION

ACTIVE INGREDIENTS

Dimethyl Phosphate of Alpha-Methylbenzyl
3-hydroxy-cis-crotonate

29.5

Petroleum hydrocarbons

59.7

INERT INGREDIENTS

14.8

Total

100.00

NET CONTENTS

Manufactured By

Economy Products Co., Inc.

SHENANDOAH, IOWA

BEST AVAILABLE COPY

5-15-67
3770-15

12/1

with conventional hand or power sprayers.
Improved control may be achieved by the addition of 1 level table
spoon of sugar (0.1 ounce) per pint of spray solution.
**For control of lice and winter ticks on dairy cattle, beef cattle,
sheep, goats and hogs.**
Mix 1 - 2 1/2 pints of CLODRIN Emulsible Concentrate (2 lb. gal.)
per 30 gallons of water (0.1% - 0.25% concentration). Spray
animals thoroughly using up to 2 gallons of spray for the 0.1%
concentration and up to 1 gallon of spray for the 0.25% con-
centration. Apply a second application 14 days later. Repeat ap-
plications as necessary to maintain control.

PRECAUTIONS IN USING

Do not contaminate feed or foodstuffs. To protect fish and wildlife,
do not contaminate streams, lakes or ponds with this material.
During commercial or prolonged exposure in spray-mixing and
loading operations wear clean synthetic rubber gloves and a mask
or respirator of a type found suitable by the U.S. Department of
Agriculture for protection against CLODRIN Insecticide. Use this
material only for recommended purposes and at recommended
dosages.

**WARNING! HAZARDOUS IF SWALLOWED, INHALED, OR AB-
SORBED THROUGH SKIN.**

Wash thoroughly with soap and water after handling and before
eating and smoking. Wear clean clothing. In case of spillage on
clothing or clothing, immediately remove clothing and flush skin or
eyes with plenty of water. For eyes, get medical attention. Keep
away from heat and open flame.

NOTICE OF WARRANTY

Economy Products Co., Inc. warrants its liability being limited to
the purchase price of the product named on this label; that the
product consists of the ingredients specified but makes no other
warranty or representation of any kind, expressed or implied, in-
cluding any warranty or representation regarding the effect or re-
sult of this product's use, whether or not the product is used in
accordance with directions. ~~AMT~~ shall have no responsibility what-
soever for injury to persons or loss or damage to property re-
sulting from the handling, storage or use of this product. The user
or buyer shall be deemed to have accepted these conditions, which
may be varied only by agreement in writing signed by duly
authorized representative of the above-named company.

BEST AVAILABLE COPY

CIODRIN® Insecticide Emulsible Concentrate is suitable for use in conventional hydraulic sprayers or low pressure knapsack sprayers. Just prior to use, mix the required amount of CIODRIN® Insecticide Emulsible Concentrate in sufficient water and apply in accordance with State and Federal Recommendations.

For control of stable flies, horn flies, face flies, house flies, lice and lone star ticks on dairy cattle, beef cattle, sheep, goats and hogs.

DIRECTIONS FOR LIVESTOCK PEST CONTROL

Mix 1 quart of CIODRIN® Emulsible Concentrate (2 lb/gal) per 6 gallons of water (1% concentration). Spray to thoroughly cover all parts of the animal, including the legs, using 1 - 2 pints of spray per large animal and proportionately less for smaller animals.

For pen or corral operations involving spray applications to livestock with high pressure equipment, do not exceed a 0.5% concentration. Mix 1 quart of CIODRIN® Emulsible Concentrate (2 lb/gal) per 12 gallons of water (0.5% concentration). Spray to thoroughly cover the animal's body and legs (avoid excessive spraying of the head) using 1 - 2 quarts of spray per large animal and proportionately less for smaller animals.

Improved face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6 - 10 gallons of spray. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

Low volume application for control of face fly

Mix 2 tablespoons (1 fluid ounce) of CIODRIN® Emulsible Concentrate (2 lb/gal) per 1½ pints of water (1% concentration). Using calibrated sprayer, apply 5 ml. of spray to the face and a total of 5 - 10 ml. to the back and sides of the animal 3 times per week.

For control of house fly, face fly and stable fly in and around livestock buildings.

Mix 1 quart of CIODRIN® Emulsible Concentrate (2 lb/gal) per 6 gallons of water (1% concentration). Spray premises such as inside and outside barn walls (including dairy barns), eaves, fences and other resting or breeding places at least twice monthly using approximately 1 gallon of finished spray per 1000 sq. ft.

BACK-RUBBER APPLICATIONS

For use in back-rubbers or face-rubbers, for the control of horn flies and face flies, mix 1 quart of CIODRIN® Emulsible Concentrate (2 lb/gal) per 7 gallons of back-rubber base oil (1% concentration). This may be diesel fuel or combinations of diesel fuel and motor oil in accordance with local recommendations.

USDA REG. NO. 3770-15



CIO

CAUTION: (Cont)
KEEPING:
ACTIVE IN
Dimet
3-hydr
Petroleum
INERT INC

Econ

4-
5-15-
3770-