

**GENERAL DIRECTIONS**

A specially prepared ready-to-use insecticide suitable for use in conventional hand, power or automatic spray equipment and in back rubbers. Approved for lactating dairy cattle. No feed-off period necessary when used on beef cattle. Apply with hand, power or automatic equipment to moisten the hair only. Do not wet the skin due to possible skin irritation. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

**USE AND DIRECTIONS FOR APPLICATION**

**Lactating Dairy and Beef Cattle: Flies (Horn Fly, Stable Fly, Face Fly, House Fly)** Make a light application of Clodrin-Vapona Insecticide daily as a fine mist to thoroughly cover all parts of the animal, including the legs. Do not apply in excess of 2 fluid ounces daily per animal.

**ACCCE**  
**AUG 2**  
 Under the Federal Fungicide, and Pesticide Act, registered under EPA Reg. No.

Ready-to-Use

**CIODRIN® - VAPONA® INSECTICIDE**  
**ANIMAL SPRAY SOLUTION**

Controls Horn Flies, Stable Flies, Face Flies, House Flies and Lice  
**FOR DAIRY AND BEEF CATTLE**

ACTIVE INGREDIENTS:	% BY WEIGHT
Dimethyl Phosphate of Alpha-Methylbenzyl 3-Hydroxy-Cis-Crotonate*	1.00
2,2-Dichlorovinyl Dimethyl Phosphate	0.23**
Related Compounds	0.02**
Petroleum Oil	98.50
<b>INERT INGREDIENTS:</b>	<b>0.25</b>
<b>Total</b>	<b>100.00%</b>

\*CIODRIN® Insecticide @Shell Chemical Co.  
 \*\*Equivalent to 0.25% VAPONA® Insecticide

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. Avoid inhalation of mist and contact with skin. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. If swallowed, induce vomiting immediately and get medical attention.

Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not reuse container. Destroy when empty.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME**

NET CONTENTS 1 GALLON

Manufactured by  
**ECONOMY PRODUCTS COMPANY, INC.**  
 SHENANDOAH, IOWA 51601

**PTED**  
**2 1975**  
 Insecticide, Fungicide, and Pesticide  
 3770-29

**USE AND DIRECTIONS FOR APPLICATION**

**Lactating Dairy and Beef Cattle: Lice** Make 1-2 applications of 3-3½ ounces of Clodrin-Vapona Insecticide at 14 day intervals as a fine mist spray to thoroughly cover all parts of the animal. If material is applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

**Back and Face Rubbers:** For the control of horn flies and face flies on lactating dairy and beef cattle, use Clodrin-Vapona Insecticide in back rubbers and face rubbers. Use 1 gallon for each 20 linear feet of burlap back rubber. For automatic back rubbers recharge the reservoir as needed. Cattle must rub face on rubbers for best face fly control.

**NOTICE:** Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 3770-29  
 EPA Est. 3770-IA-1  
 3770-NB-1 ←

8176B-MBEP00