



# 1% VAPONA<sup>®</sup> INSECTICIDE

## ACTIVE INGREDIENTS:

2,2-dichlorovinyl dimethyl phosphate  
Related Compounds  
Petroleum Distillate

## INERT INGREDIENTS:

## By Weight

0.93%\*

0.07%\*

97.00%

2.00%

100.00%

\*Equivalent to 1.00% w of Vapona Insecticide  
RShell Chemical Company

FOR LACTATING DAIRY CATTLE, BEEF CATTLE AND BARNS

## WARNING

KEEP OUT OF REACH OF CHILDREN

See left side panel for additional Warning Statements

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

NET CONTENTS 1 GALLON

Manufactured for:

**VALLEY SALES CO., INC.**

LEXINGTON, NEBRASKA 68850

ACCEPTED WITH COMMENTS

ACCEPTED

7-11-69

FUNGICIDE AND RODENT  
FOR ECONOMIC POISON  
ED UNDER NO

## DIRECTIONS

Apply **VAPONA Insecticide** as a mist spray with hand or automatic sprayer for control of face flies, stable flies, house flies, horn flies, gnats and mosquitoes.

### USE AS AN ANIMAL SPRAY

On dairy cows, beef cattle and horses, apply 2 fluid ounces of **VAPONA Insecticide** to each animal each day. The mist spray should be applied with a good hand or automatic sprayer. Spray to wet hair. For maximum results apply the 2 fluid ounces in 2 applications 1 ounce morning and night. Special care should be used to assure coverage of the legs, belly, shoulders and head. Caution: Do not apply regularly to calves under 6 months of age. Spray animal thoroughly but do not wet skin. Keep unprotected persons out of operating area.

In a complete fly control program this spray should be used in conjunction with a **VAPONA Insecticide** building, fog or surface spray and sanitary measures. Do not use this solution for fogging purposes within 8 hours of direct spray to animals.

### WARNING

May Be Fatal If Swallowed, Inhaled Or Absorbed Through The Skin. Do not get in eyes or on skin. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water, milk or milk handling equipment. Keep out of reach of children. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

### NOTICE

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk to persons or property arising out of use or handling and accepts the product on these conditions.

USDA Reg. No. 3779-11

1169A-MEELCO

## DIRECTIONS

**VAPONA Insecticide** is a very effective fog or surface spray and will provide fast knockdown and kill of flies, mosquitoes and other small flying insects. It is a multi-purpose formulation designed for use as an animal spray to protect dairy cows, beef cattle and horses from face flies, stable flies, house flies, horn flies, gnats and mosquitoes.

### USE AS A BARN FOG

**VAPONA Insecticide** can be used in barn fogging equipment without dilution. One pint will fog 8,000 cubic feet of enclosure. Before applying, reduce air movement as much as possible by closing doors, windows and other openings. Do not contaminate water, feed or foodstuffs, milk or milking utensils. Do not use this solution for fogging purposes in areas where animals have received direct application within 8 hours.

### ANIMAL BUILDINGS

(Horse Barns, Dairy Barns, Shelter Sheds),  
**MILK SHEDS, OUTDOOR AREAS**

Dilute **VAPONA Insecticide** with equal amounts of deodorized kerosene or fuel oil. Apply as a coarse wet spray at the rate of one quart per 1,000 square feet to walls, ceilings, stalls, feed storage rooms, alleyways, pig pens, milk sheds, loafing areas. In poultry houses apply to manure, window sills and outside of penned enclosures. For fly and maggot control in breeding areas such as dumps and manure piles apply at the rate of 1 to 2 quarts of the diluted mixture per 100 square feet.

Apply this diluted mixture as a coarse wet spray at the rate of one quart per 1,000 sq. feet to treat outdoor living areas, picnic grounds, patios, backyard areas, loading docks, outdoor latrines, parking areas, refuse areas around service stations, open air drive-ins, garbage collection and disposal areas.

ACCEPTED