CAUTION

Do not store near fertilizers, seeds, insecticides or fungicides

To not graze dairy animals on treated areas within 6 weeks after application.

Do not graze meat unimals on treated areas within 2 weeks of slaughter.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Coarse sprays are less likely to drift.) Vapors from this product may injure susceptible plants in the immediate vicinity. Thoroughly clean spray equipment with a suitable chemical cleaner hafore using for other purposes or do not use same spray equipment in applying other materials to plants. Removal of traces of 2.4-D and 2.4.5-T from wooden containers is almost impossible.

Do not use on lawns, around the home or recreation areas.

Do not contaminate irrigation ditches or water used for domestic purposes.

This product will kill fish. Keep out of any body of water. Apply this product only as specified on this label.

Rinse equipment and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes. Do not burn container as resulting vapors may damage nearby plants, crops and shrubbery.

Do not dispose of poison by burning since smoke from the burning vine is poisonous regardless of how killed.

Keep container tightly closed when not in

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 2 State

0.06 V-MBEPCO

NO. 2 & 2 BRUSH KILLER

Anteression manuscopposition of the contraction of

EACH GALLON CONTAINS 2 LBS. EACH OF 2.4-D AND 2,4,5-T ACID EQUIVALENT

ACTIVE INGREDIENTS:

Butyl Ester of 2,4-Dichlorophenoxyacetic Acid* Butyl Ester of 2,4,5-Trichlorophenoxyacetic Acid**		
·	Total	100.0%

*Equivalent to 2,4-Dichlorophenoxyacetic acid 21.7% or 2 lbs. 2,4-D acid equivalent per gallon

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May cause skin irritation. Avoid contact with eyes, skin or clothing. See side panel for additional caution statements.

DO NOT USE EMPTY CONTAINER --- PERFORATE, CRUSH, BURY DO NOT BURN

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

NET CONTENTS ____ GALLONIA CCEPTE

May 21973

Manufactured by

ECONOMY PRODUCTS CO., INC. SHENANDOAH, IOWA 51601

DIRECTIONS FOR USE

This product is intended only for the control of brush and deciduous woody plants such as tree sprouts. It is not interded for use on growing crops but is for use on power line rights-of-way, stump land, and certain waste land areas where it is desired to kill or repress the growth of woody plants. Ffective in the dosage indicated against e following shrubs and trees:

 \mathbf{Ash} Maple Aspen Oak Birch

Dewberry

Osage Orange Blackberry Poison Ivy Crab Apple Poplar

Sumac Dogwood Virginia Creeper

Elm Walnut Wild Grape Hawthorn Hickory Wild Cherry Wild Rose Honeysuckie Willow Locust

FOLIAGE TREATMENT (Late Spring) Application): Use 1 to 2 quarts (\(\frac{1}{2} \) to 1 lb. each of 2.4-D and 2.4.5-T; with sufficient water to adequately cover one acre. Apply early in the year when the new growth is succulent and rapidly growing. Repeat application in 4 to 6 weeks if needed. For application to smaller areas by hand or knapsack sprayers use 12 fluid oz, per gallon of water and apply at the rate of 1 gallon per sq. rod (272 sq. ft.)

BASAL BARK TREATMENT (Dormant or Growing Season Application): Use 1 pint ($\frac{1}{2}$ lbs. each of 2,4-I) and 2,4,5-T) with 3 gals, of diesel oil or fuel oil and spray the basal part of the stem or trunk from the ground level to a height of 12 to 24 inches. Circle the trunk completely and spray until run-off occurs. Apply at any time of year, using a knapsack sprayer or low volume equipment.

STUMP TEATMENT (Dormant or Growing Se son Application: Dilute 1 pint 2.4-D and 2.4.5-T with 3 gallons of diesel oil or fuel oil and apply with knapsack sprayer or low volume equipment. The bark as well as the cut surface should be wet thoroughly until run-off occurs. Treat freshly-cut stumps at any time, including winter months. Best results are usually obtained on stumps 1 inch across or larger. Repeat application if new growth develops

^{*} Equivalent to 2,4.5-Trichtorrphenoxyacetic acid 21.7% or 2 lbs. 2.4.5-T acid equivalent per gallon

label copy

2,4-D LAME WEED KILLER (Granular)

ACTIVE HIGREDIESES:

2,4-Dichloropheno:yacetic acid, Iso-octyl ester * IMERT INGREDIENTS:

97.59%

*2,4-Dichlorophenomyacetic acid equivalent 1.6% by weight

Net Weight(8 oz.) (1 lb.)

(18 pt. type) CAUTICAL (12 pt. type) IMEP OUT OF REACH OF CHILDREN Do not take internally. Avoid contact with eyes, skin and clothing. In case of skin contact wash thoroughly with soap and warm water. Thush eyes a with clean water and obtain prompt medical attention. Do not store near seeds, fertilizers, insecticides or fungicides. Avoid drift to susceptible plants as this product may injure cotton, beans, peas, tomatoes, grapes, ornamentals. Do not use in greenhouses or on crops. Do not contaminate water intended for irrigation or domestic use. Vapors from this product may injure susceptible plants in the immediate vininity. Do not reuse container. Destroy when empty.

DIRECTIONS

GRANULAR LAME WEED KILLER is a convenient form of 2,4-D as low volatile ester for killing certain broad-leaved weeds in lawn and turf. Enters the plant through the leaves and works downward to roots, killing roots and all.

LAWN: To kill broad-leaved weeds such as dandelions, plantain and other di-cot (broad-leaved) weeds, apply 8 ounces per 212 square feet or 1 pound par per 425 square feet of lawn area. Apply on a class day to avoid wind drift. Make application very evenly by sprinkling on the ground with shaker can with no skips or misses. It is suggested that the area be covered twice (cach coverage must be at half dose), the second time at right angles to the first to insure thorough coverage. Do not apply within three feet of shrubs, flowers, or other desirable plants. Repeat application in one month to control weeds that were skipped. If weeds are over 5 to 6 inches tell, they should be sprayed with Liquid Lawn Weed Killer. Do not use on St. Augustine grass. Bent grass. Dichondra. Carpetgrass or clover laws. Note: Do not use grass, Bent grass, Dichondra, Carpetgrass or clover laws. <u>Note:</u> Do not use on grass seedlings or newly seeded lawns. Use only on established grasses.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk to persons or property arising out of use or handling and accepts the product on these conditions.

USDA Reg. No. 3770-43

MBEPCO

Monufactured by ECONORY PRODUCTS CO., INC. Shenandoah, Iowa 51.601

19 FEB 1970