

label copy

2,4-D LAWN WEED KILLER
(Granular)

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid, iso-octyl ester * 2.41%

INERT INGREDIENTS:

* 2,4-Dichlorophenoxyacetic acid equivalent 1.6% by weight 97.59%

Net Weight(8 oz.) (1 lb.)

(10 pt. type) CAUTION

(10 pt. type) KEEP OUT OF REACH OF CHILDREN.

Do not take internally. Avoid contact with eyes, skin and clothing. In case of skin contact wash thoroughly with soap and warm water. Flush eyes with clean water and obtain prompt medical attention. Do not store near seeds, fertilizers, insecticides or fungicides. Avoid drift to susceptible plants as this product may injure cotton, beans, peas, tomatoes, grapes, ornamentals. Do not use in greenhouses or on crops. Do not contaminate water intended for irrigation or domestic use. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not reuse container. Destroy when empty.

DIRECTIONS

GRANULAR LAWN WEED KILLER is a convenient form of 2,4-D as low volatile ester for killing certain broad-leaved weeds in lawn and turf. Enters the plant through the leaves and works downward to roots, killing roots and all.

LAWN: To kill broad-leaved weeds such as dandelions, plantain and other di-cot (broad-leaved) weeds, apply 3 ounces per 212 square feet or 1 pound ~~per~~ per 425 square feet of lawn area. Apply on a clear day to avoid wind drift. Make application very evenly by sprinkling on the ground with shaker can with no skips or misses. It is suggested that the area be covered twice (each coverage must be at half dose), the second time at right angles to the first to insure thorough coverage. Do not apply within three feet of shrubs, flowers, or other desirable plants. Repeat application in one month to control weeds that were skipped. If weeds are over 5 or 6 inches tall, they should be sprayed with liquid lawn weed killer. Do not use on St. Augustine grass, Bent grass, Dichondra, Carpetgrass or clover lawns. Note: Do not use on grass seedlings or newly seeded lawns. Use only on established grasses.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk to persons or property, arising out of use or handling and accepts the product on these conditions.

U.S. Reg. No. 8770-40

123190

Manufactured by
E.I. du Pont de Nemours & Co., Inc.
Wilmington, Delaware

front panel

ECONOMY

LAWN WEED KILLER

Contains 2,4-D in Low-Volatile Esters
and
SILVEX* (2,4,5-TP)

(picture of
dandelion)

For Common Chickweed, Mouse-
Eared Chickweed, White Clover,
Black Medic & certain broad-
leaf weeds in lawns.

(Picture of
plantain weed)

ACTIVE INGREDIENTS:

**Isooctyl Ester of (2,4-dichlorophenoxyacetic acid) 9.06%
*Isooctyl Ester of Silvex (2-(2,4,5-trichlorophenoxy)
propionic acid) 4.26%

INERT INGREDIENTS:

**2,4-Dichlorophenoxyacetic acid equivalent 6.01% by weight
or 0.50 lb. per gallon.
*2-(2,4-dichlorophenoxy) propionic acid equivalent 3.11% by
weight or 0.25 lb. per gallon.

(1) SEE EYE CAUTION


(1) KEEP OUT OF REACH OF CHILDREN

See side panel for additional cautions

Contains 1/4 lb. Silvex per gallon

Net Contents One Pint

Manufactured By
ECONOMY PRODUCTS COMPANY INC.
Shenandoah, Iowa 57078



(right side panel)

CAUTION

Harmful if swallowed. Wash thoroughly after using. MAY CAUSE SKIN IRRITATION, AVOID CONTACT WITH EYES, SKIN, AND CLOTHING. DO NOT STORE WITH, OR NEAR, FERTILIZERS, SEEDS, INSECTICIDES, OR FUNGICIDES. It is almost impossible to remove residues of this product from sprayers and spray equipment, particularly from non-metallic parts, e.g., wood, rubber, polyethylene, or fibre, and the equipment used to measure and apply this weed killer should not be used subsequently for applying other materials to plants or crops.

(back panel)

DIRECTIONS FOR USE --LAWNS

To control common chickweed, mouse-eared chickweed, white clover, plantain, black medic, dandelion, buckhorn, ragweed, and lambsquarter, in established lawns and turf, use 11½ tablespoonfuls (5-3/4 fluid ounces) in 3 gallons of water and apply mixture as a spray to wet uniformly the weed plants on 1,000 sq. ft. (20' x 50') of area (equivalent to 2 gallons in 25 to 40 gallons of water per acre). APPLY ONLY IN THE FALL OR EARLY SPRING, since application during hot, dry weather may cause excessive damage to grass. Do not use on bentgrass, St. Augustine grass, Dichondra, Carpetgrass or clover lawns. Do not apply to Dichondra or Lippia lawns. Delay reseeding treated areas for

Manufactured By
ECONOMY PRODUCTS COMPANY INC.
Shenandoah, Iowa 57078

FOR ECONOMIC ACTION REGISTERED
AS UNDER NO. 123456789
TO ATTACHED COMMENTS.

(right side panel)

CAUTION

Harmful if swallowed. Wash thoroughly after using. MAY CAUSE SKIN IRRITATION, AVOID CONTACT WITH EYES, SKIN, AND CLOTHING. DO NOT STORE WITH, OR NEAR, FERTILIZERS, SEEDS, INSECTICIDES, OR FUNGICIDES. It is almost impossible to remove residues of this product from sprayers and spray equipment, particularly from non-metallic parts, e.g., wood, rubber, polyethylene, or fibre, and the equipment used to measure and apply this weed killer should not be used subsequently for applying other materials to plants or crops.

(back panel)

DIRECTIONS FOR USE --LAWNS

To control common chickweed, mouse-eared chickweed, white clover, plantain, black medic, dandelion, buckhorn, ragweed, and lamb's quarter, in established lawns and turf, use 11½ tablespoonfuls ~~KK~~ (5-3/4 fluid ounces in 3 gallons of water and apply mixture as a spray to wet uniformly the weed plants on 1,000 sq. ft. (20' x 50') of area (equivalent to 2 gallons in 25 to 40 gallons of water per acre). APPLY ONLY IN THE FALL OR EARLY SPRING, since application during hot, dry weather may cause excessive damage to grass. Do not use on bentgrass, St. Augustine grass, Dichondra, Carpetgrass or clover lawns. Do not apply to Dichondra or Lippia lawns. Delay reseeding treated areas for at least 3 weeks. To avoid damage to turf, do not apply in excess of the above stated quantity or rate. Do not overlap. Avoid double ~~XXXXX~~ coverage when ~~XXXXX~~ backing up and at turns. Do not spray onto the grass when nozzle is not moving, such as when cleaning or testing nozzles. Use a positive quick shut-off valve on spraying equipment. APPLY ONLY TO ESTABLISHED GRASS -- THOROUGHLY ROOTED AND TILLERED. Do not apply to newly seeded areas or to seedling grass.

CAUTION

Considerable caution must be exercised in using this product to avoid injury to crops and desirable plants. Do not apply to vegetables, flowers, grapes, fruit trees, ornamentals, cotton or other crops or desirable plants and do not permit spray mist to drift onto them since even minute quantities may cause severe injury during both growing and dormant periods. Apply only when there is no hazard from drift. Course sprays are less likely to drift. High temperatures during or shortly after application of the product may cause vaporization which would injure susceptible plants growing nearby. Do not apply excessive quantities. Excessive amounts in the soil may temporarily inhibit seed germination and plant growth. Do not contaminate domestic or irrigating water supplies.

MBEPCO

USDA Reg. No. 3770-48



UP	DATE
FILE	TIME
FILE NO	REGISTERED
OTHER NO	

✓