

## DIRECTIONS

Apply dust directly from can by punching out holes in top, or apply with hand duster.

### ANTS

**In Lawns.** Apply directly to anthills and water thoroughly to get material down into nests. Also apply to ant trails and surrounding areas. **In Gardens** for tomatoes, peas, beans, and cabbage, apply to the soil around base of plants. Also apply directly to plants covering both upper and under sides of leaves. Observe the **Special Cautions** under GARDEN PESTS below. **In Homes.** Apply on the ant trails and areas around door sills and window frames, openings around water pipes, heat ducts, electrical outlets, and baseboards where ants may come out in the rooms from wall spaces and partitions. Dust only areas inaccessible to children and pets.

### LAWNS

To control ants, chiggers, chinch bugs, mole crickets, ticks, apply this dust at the rate of 1 pound per 1,000 square feet. Water material down into soil thoroughly.

**Note:** Keep children and pets off treated areas until the treated areas have been well watered and have dried completely. Birds feeding on treated areas may be killed.

### SOIL INSECTS

**Gardens:** To control wireworms, white grubs, and Japanese beetle grubs attacking potatoes, corn, cucumbers, tomatoes, peas, and beans, apply at the rate of 1 pound per 500 square feet as preplanting soil application or at time of planting or transplanting and work into the top 2 to 3 inches of soil. Remains effective entire season.

### IN HOMES

Apply as directed below. Repeat applications as needed to maintain effective control.

**Ants:** See directions under Ants. **Roaches, waterbugs, silverfish.** Dust into infested cracks, under sinks, behind baseboards and known hiding and resting places of these insects. For **silverfish**, also treat shelving and bookcases. **Box elder bugs.** Apply around doorsills, window frames, or wherever these pests enter premises. Apply to outside areas and places where these insects congregate. **Crickets, spiders, centipedes.** Apply dust to dark places and crevices in basements, closets, and behind baseboards. **Carpet beetles.** Apply to area of the floor, baseboards, and closet shelves and work the dust into cracks where the insects may be found. Treat the underside of carpet. Cover or remove fish bowls and aquaria before and approximately one (1) hour after treatment.

### GARDEN PESTS

**On Flowers.** To control ants, armyworms, cucumber beetles, flea beetles, grasshoppers, climbing cutworms, leaf miners, tarnished plant bugs, thrips on

dahlias, gladiolus, roses, zinnias, other flowers, ornamentals and shrubs, apply to plants covering both upper and under sides of foliage thoroughly. Repeat applications as needed.

**On Vegetables.** Apply to plants covering both upper and under sides of foliage thoroughly. Apply before plants are infested by the following. Repeat applications as needed. To control **Colorado potato beetle, potato flea beetle,** on potatoes; **Flea beetles, grasshoppers, tomato hornworms,** on tomatoes; **leaf miners** on peppers; **cabbage worms, harlequin bugs, cabbage loopers,** on cabbage; **spittlebugs, strawberry crown borer,** on strawberries; **stinkbugs** on beans. To control **crickets, strawberry root weevil** on strawberries, apply to ground around base of plants and work into top 2 or 3 inches of soil. To aid in the control of **onion maggots** on onions, apply the dust around the base of the plants as soon as they are set in the ground, then the insecticide should be washed or stirred in the soil. Do not apply to spring or green onions.

### SPECIAL CAUTIONS

Do not apply after edible parts start to form on cucurbits, may be injurious to some varieties. Do not use on cabbage after heads or other edible portions begin to form, or tomatoes, peppers, strawberries after fruit begins to form; nor on green beans or snap beans after pods begin to form. Do not use on peas within 15 days of harvest. Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not use on leafy vegetables such as lettuce, spinach. Do not apply to crops to be fed to animals. Do not apply when weather conditions favor drift of dust from areas treated.

### CAUTION

Harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed, foodstuffs, water and cooking utensils.

**Container Disposal:** Do not reuse empty container. Destroy by punching holes in it and burying with wastes away from water supplies or by burning. Stay away from smoke or fumes.

**Environmental Cautions:** This product will kill fish, birds and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**NOTICE:** Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 3770-51  
EPA Est. 3770-NB-1

4085B-MBEP/CO

# 6% CHLORDANE DUST

Insecticide,  
Federal Insecticide, Fungicide and Rodenticide Act,  
as amended by the pesticide  
amendment under  
EPA Reg. No. 3770-51

## For The Control of

Ants - Silver Fish - Cockroaches & Certain Other Insects Infesting

## GARDENS -- LAWNS -- HOMES

### ACTIVE INGREDIENT:

Technical Chlordane\* ----- 6%

INERT INGREDIENTS: ----- 94%

\*Equivalent to 3.6% Octachloro-4,7-Methanotetrahydroindane and 2.4% related compounds.

## CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See Side Panel for Additional Cautions.

NET WEIGHT 1 POUND

Manufactured by:

ECONOMY PRODUCTS COMPANY, INC.

SHENANDOAH, IOWA 51601