

SEVIN GARDEN SPRAY

A 50% Wettable Powder for control of Japanese
Beetle, Squash bugs, certain other insects

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 50%

INERT INGREDIENTS: 50%

*Trade Mark of Union Carbide Corp. for the active ingredient,
1-naphthyl N-methylcarbamate
U. S. Pat. No. 2,903,478; 3,009,855

For ORNAMENTALS - FLOWERS - LAWNS - FRUITS - VEGETABLES

CAUTION
KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions.

NET WEIGHT 8 OZ.

Manufactured by:
ECONOMY PRODUCTS COMPANY
SHENANDOAH, IOWA 51601

DIRECTIONS

Mix 2 level tablespoonfuls Sevin Garden Spray in 1 gallon of water. To use, mix designated amount of Sevin Garden Spray in a little water to make a thin paste, then add required amount of water. Do not store diluted spray. Make new dilution for each use. Keep spray mixture well agitated.

ROSES, ORNAMENTALS, FLOWERS, SHRUBS, SHADE TREES: Japanese Beetles, Aphids, Bagworm, Elm Leaf Beetle, Certain Leafminers, Rose Slug, Periodical Cicada (17-year locust), Scale Insects, Tent Caterpillars, Blister Beetles, Leaf Rollers, Exposed Thrips. Start spraying when insects or their damage first appear. Repeat weekly or as necessary. Do not use on Boston Ivy.

FRUIT TREES — APPLES, PEARS AND PEACHES: Apple Maggot, Plum Curculio, Apple Aphid, Leaf Rollers, Lesser Peach Tree Borer, Oriental Fruit Moth, Codling Moth. Make one application between bud break and bloom, a second application at petal fall, and then at 3 to 4 week intervals for 3 to 4 months as necessary. To control rosy apple aphids apply before leaves are curled. Do not apply later than 1 day before harvest. (Note: To avoid possible fruit thinning on apples and foliar injury on McIntosh, do not apply within 15 days after petal fall.)

LAWNS: Fleas and Ants. Use 6 ounces in 15 gallons water and cover 850 sq. ft. of established lawn. For best results, use lawn sprayer. Mow lawn first and apply after rain or watering. Do not water for at least two days after application. Repeat in 2 to 3 weeks or as necessary.

GARDEN VEGETABLES

Use 2 tablespoons per gallon of water unless otherwise specified. Use sufficient spray to obtain full coverage. Keep spray mixture well agitated. Apply when insects or their damage appear. Repeat at 7 to 10-day intervals or as necessary.

POTATO, TOMATO, EGGPLANT and PEPPERS: Use 2 tablespoons per gallon of water for Colorado potato beetle, flea beetles and leafhoppers. Use 2 to 4 tablespoons per gallon for European Corn Borer, fall plant bugs and stink bugs.

CUCUMBERS, MELONS, PUMPKINS and SQUASH: Pickleworm and melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive application. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

SWEET CORN: Use 2 to 4 tablespoons per gallon of water for corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetle, sap beetles and leafhoppers.

For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary.

For insects attacking silks and ears, apply at 2 to 3-day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

CAUTION: Do not breathe dust. Do not take internally. Harmful if swallowed or inhaled. Avoid breathing dust or spray mist. Avoid contact with skin; wash thoroughly after using. Avoid storage near feed and food products. Keep children and pets off treated areas until this material is washed into the soil and grass is dry. Do not reuse container. Destroy when empty.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 3770-55

SEVIN-MBEP-60