

MIXING DIRECTIONS

BEEF CATTLE and HOGS: For control of Horn Flies, Lone Star Ticks, Winter Ticks, Gulf Coast Ticks, Spinose Ear Ticks, Lice and Sarcoptic Mange Mites—Mix 1 gallon of **Range Cattle Spray Conc.** in 100 gallons of water (2 1/2 tablespoonfuls to 1 gallon of water) for either spraying or dipping.

SHEEP and GOATS: For control of Sheep Ticks (Keds) and Lice—Mix 1 quart of **Range Cattle Spray Conc.** in 125 gallons of water for spraying; mix 1 quart of **Range Cattle Spray Conc.** in 250 gallons of water for dipping vat solution. For the control of Scabies Mites—Mix 1 gallon **Range Cattle Spray Conc.** in 100 gallons of water used in dipping vat.

NOTE: Do not dip animals within 60 days of slaughter. Do not spray animals within 30 days of slaughter. Do not apply to dairy animals or animals under 3 months of age.

USE DIRECTIONS

Mix spray or dipping vat solutions thoroughly before using and keep well mixed while spraying or dipping animals. Repeat treatment at monthly intervals, if necessary. Do not spray animals confined in closed buildings. Do not treat horses or mules. Do not treat weak or emaciated animals.

CATTLE: Use sufficient pressure to wet the animals to the skin. For Gulf Coast Ticks and Spinose Ear Ticks, apply at low pressure to each ear and to the head, neck and shoulders.

SHEEP and GOATS: Apply 2 to 4 quarts of spray solution per animal at low pressure to lambs and adults after spring shearing. Use high pressure to penetrate fleece during fall treatment. (When dipping make sure the animal is held in the vat at least two minutes and submerge each animal twice during dipping).

HOGS: Spray or dip entire animals so as to thoroughly wet the skin, including the underside. Do not treat sows within 30 days of farrowing nor for at least 3 weeks thereafter. Avoid treating animals in cold, stormy weather. Do not treat pigs before weaning.

CATTLE RUBBING DEVICES

To aid in controlling Horn Flies, Mosquitoes, Gnats and Lice on **BEEF CATTLE**—thoroughly mix together 1 pint of **Range Cattle Spray Conc.** with 1 gallon of No. 2 fuel oil or diesel fuel oil in a separate container. (Do not mix with used motor oil, crankcase drainings or greases. Do not use in automatic oilers which allow an excessive or irregular amount of oil on the rubbing surface.) Apply the mixed solution to the burlap covering or pour into oil reservoir until canvas covering is saturated. Replenish the units with suitable amounts of the solution every 2 to 3 weeks. Regular exposure of the animals for about 10 weeks is usually necessary for good results against lice. Do not expect complete control of lice. **Note:** Do not allow calves under 6 months of age, or dairy animals to treat themselves. Do not permit meat animals access to backrubbers within 30 days of slaughter.

Economy

L - T PLUS

RANGE CATTLE SPRAY CONC.

FOR SPRAYING OR DIPPING VATS

Kills Horn Flies, Ticks, Lice, Mosquitoes, Gnats,
Psoroptic Scabies and Sarcoptic Mange Mites.
(Agricultural Use — Grasshopper Control)

ACTIVE INGREDIENTS:

*Toxaphene	45.00%
Gamma Isomer of Benzene Hexachloride (from Lindane)	0.45%
Aromatic Petroleum Derivative Solvent	41.55%
Xylene	5.00%

INERT INGREDIENTS:

.....	8.00%
100.00%	

*Technical chlorinated camphene, containing 67% to 69% Chlorine

WARNING

KEEP OUT OF REACH OF CHILDREN
See right side panel for additional cautions.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

NET CONTENTS

Manufactured by

Economy Products Co., Inc.

SHENANDOAH, IOWA 51601

FOR GRASSHOPPER CONTROL

Spray or bait hatching areas such as roadsides, field margins or idle land as soon as grasshoppers are seen.

For Spraying: Use 1 quart of **Range Cattle Spray Conc.** per acre in sufficient water to provide uniform coverage for newly hatched grasshoppers and 1 1/2 quarts Conc. per acre for mature grasshoppers. Apply required amount of **Range Cattle Spray Conc.** in the quantity of water that the sprayer delivers per acre. The amount of water required per acre will depend upon the type of equipment being used. This varies from 5 to 100 gallons of water per acre.

Wet Bait: Mix 1 quart of **Range Cattle Spray Conc.** in 10 gallons of water. Apply solution to mixture of 50 pounds of wheat bran and 2 1/2 bushels of sawdust. Spread wet bait at the rate of 20 pounds per acre.

Dry Bait: Mix 1 quart of **Range Cattle Spray Conc.** with 1 quart of kerosene. Apply oil solution as a fine spray to 100 pounds of coarse bran. Spread dry bait at the rate of 5 pounds per acre.

WARNING

May be fatal if swallowed! Do not breathe spray mist. May be absorbed through skin. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Do not store near feed and food products.

Do not use on dairy animals or in dairy barns, or in poultry houses. Avoid contamination of feed, foodstuffs, feeders, waterers and food equipment. Do not feed treated forage to dairy animals or animals being finished for slaughter.

This product will kill fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not reuse container. Destroy when empty.

NOTICE: Sellers guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

USDA Reg. No. 3770-70

2116A-MBPCO

