

WARNING: Harmful if swallowed, inhaled or absorbed through the skin. Avoid inhalation. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. Do not allow the spray mist to contaminate milk or milking utensils. Cover feed and feed troughs or remove before spraying.

This product is an organic phosphate which is a cholinesterase inhibitor and if poisoning should occur, CALL A PHYSICIAN. Atropine is antidotal.

This product is toxic to fish and wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not reuse container. Destroy it by perforating or crushing. Bury or discard in a safe place.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

CIODRIN® DAIRY BACKRUBBER

A ready-to-use insecticide for the control of horn flies, stable flies, and face flies on lactating dairy cattle and beef cattle.

ACTIVE INGREDIENTS:

Dimethyl phosphate of alpha-methylbenzyl 3-hydroxy-cis-crotonate* 1.00%
Petroleum hydrocarbons 98.82%

INERT INGREDIENTS:

.18%
100.00%

*Ciodrin Insecticide

©Shell Chemical Company

ACCEPTED
May 1, 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTER-
ED UNDER NO. 3770-80 SUBJECT
TO ATTACHED COMMENTS.

WARNING

KEEP OUT OF REACH OF CHILDREN

See side panel for additional warning statements.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

NET CONTENTS 1 GALLON

Manufactured by

ECONOMY PRODUCTS COMPANY, INC.
SHENANDOAH, IOWA 51601

BACKRUBBER AND FACERUBBERS

DAIRY CATTLE, BEEF CATTLE - Horn flies and face flies are only controlled when the animal rubs its face on the rubber. Use undiluted in backrubber or face rubber reservoir. Recharge as necessary. If backrubber has no reservoir, saturate cable at the rate of 2 to 3 quarts per 20 feet of cable. Repeat every two weeks or more often if conditions warrant. Backrubbers or facerubbers should be located around the pasture and/or loafing areas especially near feed bunks, watering facilities, or salt licks where the cattle usually congregate so that it will be used frequently.

AS A SPRAY

For the control of face flies, horn flies and stable flies, apply as a mist spray to animals by use of an automatic sprayer (treadle or electric eye actuated) or by hand-operated sprayers at milking time. Apply at the rate of a maximum of two ounces per animal per day. For best fly control apply daily.

EPA Reg. No. 3770-80

8297A-MBEP(0)