

SEVIN* 5% DUST INSECTICIDE

FOR THE CONTROL OF INSECT PESTS ON VEGETABLES,
STRAWBERRIES, ORNAMENTALS, AND CATS AND DOGS

Controls:

Mexican bean beetles, Japanese beetles, squash bugs, fleas,
and certain other insect pests

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) .. 5.0%

INERT INGREDIENTS: 95.0%

Total 100.0%

*Trade Mark Union Carbide Corporation for 1-naphthyl N-methylcarbamate U. S. Patent 2,903,478

CAUTION

KEEP OUT OF REACH OF CHILDREN


See back panel for additional cautions

NET WEIGHT 1 POUND

Manufactured for:

GURNEY SEED & NURSERY CO.

YANKTON, SO. DAKOTA 57078



DIRECTIONS FOR USE VEGETABLES & STRAWBERRIES

Make application in morning or evening when it is cool and air is calm. Treat upper and lower surfaces of all foliage. Do not apply in excess. Start application at first sign of insects. Repeat as necessary, generally at 7-day intervals, oftener in rainy weather. Do not use closer to harvest than indicated following each crop. Light application should be made in seed beds if plants show signs of insects.

TIME TO STOP USE BEFORE HARVEST:

FOR CONTROL OF:

Asparagus: asparagus beetle, Apache cicada (do not repeat application within 8 days)1 day
Beans (green beans, lima beans, snap beans, cowpeas, blackeyed peas): Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetle, Japanese beetle, leafhopper, velvet bean caterpillar, western bean cutworm, armyworm, corn earworm, Lima bean pod borer, stink bugs and tarnished plant bugsno limit
Sweet corn: corn earworm (Apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals as infestation warrants. Re-apply immediately after rain.), corn rootworm adult, European corn borer, fall armyworm, flea beetles, Japanese beetles, sap beetles and leafhoppers (application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations)no limit
Cucumber, Melons, Pumpkin, Squash: pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bug (avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity)no limit
Potato, Tomato, Eggplant, Peppers, Peas: Colorado potato beetle, flea beetles, leafhopper, European corn borer, fall armyworm, lace bugs, tomato fruit worm, tomato hornworm, tarnished plant bugs, stink bugs:no limit
Okra: corn earworm, stink bugsno limit
Carrots: Leafhoppers, flea beetles and spittlebugsno limit
Horseradish, Parsnips, Radish, Rutabagas: flea beetle3 days
Chinese Cabbage, Collards, Dandelion, Endive (escarole), Garden Beets, Kale, Lettuce, Mustard Greens, Parsley, Salsify, Spinach, Swiss Chard, Turnips: flea beetle, harlequin bug, leafhopper, armyworms, cabbage caterpillars, corn earworm, spittlebugs, tarnished plant bug, stink bug14 days
Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi: flea beetle, harlequin bug, armyworm, cabbage caterpillar, corn earworm3 days
Strawberries: Meadow spittlebug, leaf roller and strawberry weevil1 day

ORNAMENTALS

Sevin 5% Dust can be used on ornamental plants, flowers, shrubs and small trees: rose, carnation, gladiolus, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. SEVIN injures Boston ivy.

Apply to the upper and lower leaf surface of all foliage for complete coverage to control pests: apple aphid, bagworm, birch leaf miner, blister beetles, boxelder bug, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leafroller, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillar, psyllids, rose aphid, rose-slug scale insects, tent caterpillar, thorn bugs, thrips (exposed), willow leaf beetles.

Note: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment.

CATS AND DOGS

For ticks and fleas dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Do not treat kittens and puppies less than 4 weeks of age.

In and around dog and cat quarters: For ticks and fleas dust liberally in and around sleeping quarters, kennels and dog houses by means of a shaker can, or dust gun.

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Do not take internally. Avoid inhalation of dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of food, feed, feeding troughs and water receptacles.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL CAUTION: This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

CONTAINER DISPOSAL: Do not reuse container. Destroy when empty.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 3770-108-4.

EPA Est. 3770-1A-1

3770-NB-1

3495B-MBEP CO