

RATON-LYN®

Warfarin plus Sulfaquinoxaline Rodenticide

POUNDS

For rats and mice. This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent, sulfaquinoxaline, which inhibits bacteria that produce vitamin K, an antidote for warfarin.

ACTIVE INGREDIENTS:

Warfarin [3-(α -acetylbenzyl)-4-hydroxycoumarin] 0.5%

N1-(2-quinoxaliny) sulfanilamide (sulfaquinoxaline) 0.5%

INERT INGREDIENTS: 99.0%

100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

CAUTION:

Keep away from humans, domestic animals, wildlife or pets. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by over-doses of bishydroxycoumarin.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTE FOR PHYSICIANS:

When a human has been known to have ingested RATON-LYN, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol R.

NOTICE:

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

USDA Reg. No. 3770-110

DIRECTIONS FOR MIXING RAT OR MOUSE BAIT:

Mix RATON-LYN Concentrate thoroughly with cracked or ground grains, dog food, or other edible inert baiting materials. Use 1 part RATON-LYN Concentrate to 19 parts by weight of inert bait.

Place baits in locations not accessible to children, pets, wildlife and domestic animals, or in tamper proof bait boxes.

When a continuous source of infestation is present (nearby dumps) permanent bait stations should be established and replenished as needed.

DIRECTIONS FOR USE OF RAT BAIT:

Place 4 to 6 ounces of bait in dry container next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 10 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

DIRECTION FOR USE OF MOUSE BAITS:

Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations of intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of at least 15 days.

Manufactured by:

ECONOMY PRODUCTS CO.

Shenandoah, Iowa 51601

