

UNITED STATES DEPARTMENT OF AGRICULTURE
 OFFICE OF THE SECRETARY
 WASHINGTON, D. C. 20250
 RECEIVED
 JUN 10 1960
 3

FOR CONTROL OF HORN FLIES ON DAIRY CATTLE AND HORN FLIES, LICE AND TICKS ON BEEF CATTLE

ACTIVE INGREDIENTS:
 *Methoxychlor, technical 5.00%
 **Malathion 4.00%
INERT INGREDIENTS: 91.00%

Equivalent to 1.17% 2,2-bis(4-methoxyphenyl)-1,1,1-trichloroethane and 0.67% other isomers and reaction products.
 O,O-dimethyl dithiophosphate of diethyl malathion.

CAUTION KEEP OUT OF REACH OF CHILDREN

Avoid breathing of dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and water troughs. Do not reuse empty container. Destroy container when empty.

This product is toxic to fish. Keep out of any body of water.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk to persons or property arising out of use or handling, and accepts the product on the label condition.

DIRECTIONS FOR USE AS A DUST BAG

For use as a dust bag hang by rope and open bottom of outside waterproof bag.

BEEF CATTLE: To aid in the control of horn flies and lice, hang where cattle tend to congregate. For **BEEF CATTLE** a good location is near salt blocks, mineral blocks, or water supply. (Do not hang dust bag directly over feed, mineral, salt or water supply). For cattle on range or pasture, hang dust bag between fixed locations such as between 2 posts, buildings or trees. Dust Bag should be hung so bottom of bag is 4 to 6' below topline of cattle.

Allow one dust bag for each 10-15 animals.

Do not use the dust bag treatment for dairy cattle. For application to dairy cattle, follow directions given under "Directions For Application By Hand."

NET WEIGHT 10 LBS

Manufactured by:

100% PURE PRODUCT

SHENANDOAH IOWA 51601

DIRECTIONS FOR APPLICATION BY HAND

Dairy and Beef Cattle: Horn Flies. Apply 2 heaping tablespoonful (2 ozs.) to back and neck of each animal. Rub the dust into the hair. Repeat every 3 weeks if necessary. Do not apply to lactating Dairy Cattle within 5 hours of milking or during milking. When dusting dairy cattle, treatments should be made no closer than 3 weeks apart.

Beef Cattle: Horn Flies, Lice and Ticks. Make thorough application to all parts of the animal's body by hand or power duster. Work the dust down to the skin with the hand. Repeat only as necessary.

Hogs: Lice. Make thorough application to all 55% of the animal's body by hand or power duster. Also treat pen and litter. Repeat as necessary.

Sheep and Goats: Lice and Keds. Make thorough application to animals with hand or power duster. Rub dust into the wool or hair. Do not apply to lactating milk goats.