

## DIRECTIONS

**BEEF CATTLE AND HORSES:** Ticks, Lice, Horn Flies: To control Lone Star Tick, Cattle Fever Tick, Ear Tick, Cattle Lice and Horn Flies, dilute 8 ounces in 6 gallons of water (1 quart in 25 gallons of water). Dip or apply to animal body thoroughly with enough spray force to part their hair and wet the skin. If reinfestations occur repeat application in 15 to 21 days or as needed for control. Do not add infested cattle to the treated herd. **WARNING:** Do not dip calves under six months old nor apply in any way to animals under three months old. Observe 30 day waiting period between last application and slaughter of animals.

**SHEEP AND GOATS:** Lice, Sheep Ticks (Keds): Mix 8 ounces with 12 to 25 gallons of water as local recommendations indicate. Dip or spray thoroughly, preferable in spring or summer when wool is short to give best spray penetration to the skin. Replacement ewes or rams should be treated before adding to the flock. For Sheep Scabies mix 1 quart with 25 gallons of water. Dip animals making sure that material penetrates wool especially around head. A single dipping is usually sufficient. If dipping vat is intermittently used, agitate before next dipping. **WARNING:** Do not use on dairy goats or in dairy barns. Do not dip individuals under six months old nor spray animals under three months old. Do not dip within 60 days of slaughter. Do not spray within 30 days of slaughter.

**HOGS:** Lice: Use 8 ounces with 6 gallons of water (1 quart in 25 gallons of water). A single spray application will give control. Spray sows 2 weeks before farrowing and not for 3 weeks thereafter. Pigs should not be sprayed when less than weaning age. **WARNING:** Do not treat hogs within 30 days of slaughter.

**UTILITY SPRAY:** for Mosquitoes, Fleas, Gnats: To control these pests in Out-Buildings, Sheds, Barns, Fences and Waste Foliage dilute 1 quart with 10 gallons of water. Make application to Walls, Doors, Ground Areas, Ceilings and places where insects rest or breed. Repeat applications as needed for control. For fleas treat animals' bedding and replace. **WARNING:** Do not use in dairy barns, milk houses or poultry houses. Do not use the utility spray dilution on any animals. Not for use in dwellings.

**SARCOPTIC MANGE MITE:** On Hogs, Beef Cattle and Horses mix 8 ounces with 10 gallons of water (1 gallon with 150 gallons of water). Apply to herd when mange is found. In light cases a single treatment may be all that is needed. If condition persists, repeat treatment at 10-14 day intervals as needed. **WARNING:** Do not spray animals within 30 days of slaughter or do not dip within 60 days of slaughter.

EPA Reg. No. 3770-119

2200B-MBEPCC

# EPCO

## SPRAY OR DIP TO CONTROL LICE, TICKS, FLIES, SARCOPTIC MANGE MITES, FLEAS

For Sarcoptic Mange Mite, 1 gallon makes 150 gallons spray. See side panels for additional uses or additional rates.

TAKE TIME



OBSERVE LABEL DIRECTIONS

### ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene, Chlorine Content 67-69%)	45.00%
Gamma Isomer of Benzene Hexachloride (from Lindane)	1.96%
Xylene	5.00%
Petroleum distillate	35.54%
INERT INGREDIENTS:	12.50%

## WARNING

KEEP OUT OF REACH OF CHILDREN  
See side panel for additional warnings.

NET CONTENTS

JAN 26 1976

ECONOMY PRODUCTS CO., INC.  
SHENANDOAH, IOWA

Manufactured by Economy Products Co., Inc. Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered under EPA Reg. No. 3770-119

## DIRECTIONS

**FACE FLIES:** For use against face flies, one end of the back-rubber must be low enough (not more than 15 to 20 inches above ground) to allow cattle to rub their heads. The product is only effective for those cattle that rub the face and head. Dilute as for back-rubber use.

**BACKRUBBER USE:** To aid in reducing face flies and lice on beef cattle and to control hornflies, dilute 1 quart with 11 quarts fuel oil. Apply the diluted material to saturate the backrubber mechanism. Recharge the backrubber when necessary. Locate the back-rubber to give the cattle free access to it. **WARNING:** Do not allow calves under six months old or lactating dairy animals to treat themselves. Do not permit animals access to treatment within 30 days of slaughter.

**WARNING:** May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or clothing. Wash thoroughly after using. In case of contact with eyes or skin wash immediately; for eyes, get medical attention. Do not contaminate feed or foodstuffs. Do not use on dogs, cats or other household pets nor on poultry or humans under any circumstances. Not for use or storage in or around the house.

Use only recommended rate of dilution. Do not use on dairy animals. Do not use in dairy barns, milk houses or poultry houses. Do not use on food or forage to be fed to dairy animals or animals being finished for slaughter. Dip only healthy adult animals under normal conditions. Water animals before dipping. Avoid spraying or dipping animals in cold stormy weather and avoid treating overheated or sick animals. Do not spray animals within 30 days of slaughter nor dip within 60 days of slaughter. **(DO NOT USE THIS PRODUCT ON ANIMALS SIMULTANEOUSLY OR WITHIN 2 WEEKS BEFORE OR AFTER TREATMENT WITH OR EXPOSURE TO OTHER PESTICIDES OR CHEMICALS.)**

Toxic to fish and wildlife, do not contaminate streams, lakes or ponds with this material. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and container.

Do not use empty containers. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies.

Protect from Temperatures Below 32° F.

Do Not Use, Pour, Spill or Store Near Heat or Open Flame.

**WARNING: IN CASE OF POISONING CONSULT PHYSICIAN.**

**NOTICE:** Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

ACCEPTED WITH COMMENTS