

DIRECTIONS

To protect dairy stock from hornflies, houseflies, stableflies, gnats and mosquitoes, spray after cattle enter the barn, using 1 to 2 ounces per head, not to exceed 2 ounces per animal per day. The spray should be applied as a light mist primarily to reach and hit the flies as they attempt to annoy the cows.

For control of houseflies, spray with a fine mist to the ceiling areas, not over 2000 feet in barns, and use a stiff brush on stall areas to get the spray to the crevices of the stall. Treatment for flies should not be repeated more often than once every three weeks.

Space treatment in barns, houses and milk parlors for effective control of flies. Direct mist toward the upper portions of the enclosure, filling the air with fine spray particles, using approximately 2 ounces of material for each 1000 cubic feet of space. Whenever possible, keep doors and windows closed for 10 minutes after application.

Face fly spray—Apply to face of animal in morning while stalled or just before releasing to pasture. Use a spray which produces large wetting droplets. Apply sufficient material to wet the face of the animal, but do not exceed 1 1/2 ounces per animal. Repeat daily, or as necessary.

NOTICE

Seller's guarantee shall be limited to the terms of the label and subject them to the buyer, assuming any risk for persons not properly using, or for use or handling in accordance with the instructions on the label.

PYRENONE-TYPE DAIRY SPRAY

For the Protection of Dairy Animals from biting flies and the control of Houseflies in Barns and Milk Houses.

Protects against Hornflies, Stableflies, Houseflies, Gnats, Mosquitoes and Blood Sucking Lice.

A temporary treatment to relieve cattle from annoyance caused by face flies.

ACTIVE INGREDIENTS	100%
Pyrethrins	0.1%
Technical Piperonyl Butoxide**	1.0%
Mineral Oil	98.9%

**Equivalent to 0.80% of (butylcarbityl) (6-propylpiperonyl) ether and 0.20% of related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed.

Avoid contamination of milking equipment and utensils.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

NET CONTENTS 1 GALLON

Manufactured by

ECONOMY PRODUCTS COMPANY, INC.

SIEMANSDALE, IOWA 51601

DIRECTIONS

To protect dairy stock from hornflies, houseflies, stableflies, gnats and mosquitoes, spray after cattle enter the barn, using 1 to 2 ounces per head, not to exceed 2 ounces per animal per day. The spray should be applied as a light mist primarily to reach and hit the flies as they attempt to annoy the cows.

For control of houseflies, spray with a fine mist to the ceiling areas, not over 2000 feet in barns, and use a stiff brush on stall areas to get the spray to the crevices of the stall. Treatment for flies should not be repeated more often than once every three weeks.

Space treatment in barns, houses and milk parlors for effective control of flies. Direct mist toward the upper portions of the enclosure, filling the air with fine spray particles, using approximately 2 ounces of material for each 1000 cubic feet of space. Whenever possible, keep doors and windows closed for 10 minutes after application.

Face fly spray—Apply to face of animal in morning while stalled or just before releasing to pasture. Use a spray which produces large wetting droplets. Apply sufficient material to wet the face of the animal, but do not exceed 1 1/2 ounces per animal. Repeat daily, or as necessary.

NOTICE

Seller's guarantee shall be limited to the terms of the label and subject them to the buyer, assuming any risk for persons not properly using, or for use or handling in accordance with the instructions on the label.