DIRECTIONS FOR USE Dairy, Beef Cattle LIVESTOCK FLY CONTROL

FOR MIST SPRAYING USING BASE OIL: For the control of Face Flies, Horn Flies, and Stable Flies, mix 1 quart Ciodrin - Vapona Concentrate with 18 quarts base oil (diesel or fuel oil). Apply as a fine mist at the rate of 2 oz. per animal. Cover all parts of animal thoroughly. Do not exceed 2 oz. per animal per day . . . Finished spray will contain 1% Ciodrin and .05% Vapona.

MIST SPRAY—Mix 1 gallon of Ciodrin - Vapona Concentrate with 7 gallons (1 quart per 13/4 gallons) of water. Apply as a fine mist at the rate of 2 oz. per animal. Cover all parts of the animal thoroughly. Do not exceed 2 oz. per animal per day.

"POUR-ON"—For the control of Horn Flies dilute 1 gallon Ciodrin - Vapona Concentrate with 14 gallons (1 cupful per 7 pints) of water, and apply by pouring 1 quart of the diluted emulsion along the backline of the animal. Repeat as necessary but not more often than once weekly.

BACKRUBBER—For the control of Horn Flies on Dairy and Beef cattle, dilute 1 gallon of Ciodrin - Vapona Concentrate with 18 gallons of fuel oil or other suitable carrier and charge backrubber device or sack or cloth as occasion requires. Retreat backrubber as needed. Do not spray animals with this oil solution.

LICE and TICKS—For the control of Lice and Ticks on Cattle, Sheep, Goats, Horses and Hogs dilute 1 gallon of Ciodrin - Vapona Concentrate with 50 gallons of water and spray animal thoroughly. Apply one to two quarts per animal for adult cows and one quart per animal for adult sheep, goats, and hogs. Dosages should be proportionally reduced for smaller animals. Make second application in 14 days if needed. Repeat as necessary but not more often than once weekly.



Residual <u>Dairy Livestock</u> Spray and Backrubber Concentrate CONTROLS: Stable Flies, Horn Flies, Face Flies, Lice, Ticks.

WARNING Keep Out Of Reach of Children

See back panel for additional warning statement.
KEEP AWAY FROM HEAT AND OPEN FLAME

ACTIVE INGREDIENTS: dimethyl phosphate of alpha-methylbenzyl 3-hydroxy	
cis-crotonate	15.00%*
2,2-dichlorovinyl dimethyl phosphate	1.00% **
related compounds	.07 % **
aromatic petroleum hydrocarbons	71.18%
INERT INGREDIENTS:	12.75%
® Shell Chemical Co. *Ciodrin® Insecticide. **Equivale	ent to 1.07%
Vapona® Inse : 'e.	

DIRECTIONS FOR USE

BARN FLY CONTROL—As an aid in the reduction of House Flies, Face Flies and Stable Flies in Dairy barns, animal shelters, has barns and poultry-houses where Flies are a problem, mix 1 gallon of Ciodrin - Vapona Concentrate with 25 gallons (1 pint per 3 gallons) of water. Spray mixture on surfaces where flies alight and feed. Spray surface to run-off. To improve control add 1 pound of sugar to 6 gallons of spray solution.

WARNING: Hazardous if swallowed, inhaled or absorbed through skin! Do not breathe spray mist. Do not get in eyes, on skin, or clothing. Wash with soap and water after using and before eating and smoking. Avoid contamination of feed, water and foodstuffs. During commercial or prolonged exposure in spray mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type found suitable by the U. S. Department of Agriculture for protection against Ciodrin Insecticide.

To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material.

Do not use this material on Brahman or Brahman cross cattle.

Do not reuse container. Destroy when empty.

ANTIDOTE: External—In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention.

Internal—Call a physician immediately, give emetic of mustard. Large doses of Atropine are antidotal.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk to persons or property arising out of use or handling and accepts the product on these conditions.

USDA Reg. No. 3770-163

NET CONTENTS 1 GALLON





