Crabgrass Preventer

ACTIVE	By Weight			
			oterephthalic acid	
8	,	•	Total	يسه يعشو جسه



Destr buted to

Agri-Chemicals, Inc.

MEAD SOLIOWING CAREFULLY REPORT APPLICATION

LISE THIS PRODUCT ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED DOSAGE

This product is designed for the pre-emergence control of annual weeds such as smooth and hairy crabgrass, green and yellow factail, common chickweed, Florida pussley, lembaquarters, pursione and witch grass in lawns. When used as recommended, it will not harm established grass and will control above weeds for an entire season by killing plants as seeds germinate.

DIRECTIONS FOR APPLYING

WISSE TO APPLY: This product must be applied in early spring before above weeds germinate. With the exception of common chickwood, these weeds may germinate any time there is damp rainy weather with night temperatures above 65° F. for 3 to 6 days.

Stations: The ideal time to sow lawn seed is in the fall. If spring seeding is necessary, sow seed sufficiently early so there is a noticeable green stand of grass before deadline for applying the pre-emergence control. As an alternative, apply this product as soon as the frost is out of the ground and delay seeding for approximately 40 days. The shorter the waiting period, the more grass seed must be sown to incore a desirable stand. Delay of less than 30 days is not recommended.

COMMON CHICKWEED: This weed thrives in coal weather. In some areas, it is a problem in the fall and spring; in others, depending on the weather, it grows anytime. To control this weed, it is suggested that you use this product to prevent new plants from developing for a season.

HOW TO APPLY: Use a fertilizer spreader per gu'de shown below:

SPREADER SETTING GUIDE

Spreader settings are approximate and are based an an average walking speed. Reduce setting for slower speed; reise setting for higher speed.

Mate	•		,	* .	
Wards 1091, 1093, 1095				9	
Wards 1080, 1081 Central (Recessed Wheel)	*	••	 **	**	
Jeckson & Seers					
Cast					

CAUTION

Do not take internally. Avoid inholation or contact with eyes. Do not conteminate feed or foodstuffs. Week after handling or using. Do not use on dichendra.

NOTICE

UBS Agis-Chemicals, inc. can accept no responsibility for injury to parties or animals or