

(LABEL COPY)

(front panel)

FORE^R
A Fungicide for Turf and Ornamentals

ACTIVE INGREDIENTS:

| | |
|---|------|
| A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate | 80% |
| In which the ingredients are | |
| Manganese + + | 16% |
| Zinc + + | 2% |
| Ethylene bisdithiocarbamate ion | |
| (C ₄ H ₆ N ₂ S ₄) = | 62% |
| INERT INGREDIENT | 20% |
| Total | 100% |


U.S. Patents 2,504,404; 2,974,156 and Patent Pending
(R) Trademark Rohm & Haas

(18 pt. type) CAUTION

(12 pt. type) KEEP OUT OF REACH OF CHILDREN

May cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray mist. In case of skin contact wash with water; for eyes wash copiously with water and get medical attention. Contamination of fish-bearing waters with this product may harm fish.

Important: Keep away from fire and sparks. Store in a cool, dry place. Do not allow to become wet or overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness of FORE and may also generate flammable vapors. Keep container closed when not in use.



(back panel)

Net Contents (1 lb.) (5 lbs.)

Repacked by
ECONOMY PRODUCTS COMPANY INC.
Shenandoah, Iowa 51601

GENERAL INFORMATION


To mix, slowly sift FORE Turf and Ornamental Fungicide into spray tank ~~and~~ or container while it is being filled, keeping solution well agitated or stirred. Add insecticide and nutritional sprays last. FORE Turf and Ornamental Fungicide is believed compatible with most commonly used pesticides.

DIRECTIONS FOR USE

TURF

Fusarium Blight (F. roseum), Red Thread, Copper Spot, Slime Molds--In a

Important: Keep away from fire and sparks. Store in a cool, dry place. Do not allow to become wet or overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness of FORE and may also generate flammable vapors. Keep container closed when not in use.



Net Contents (1 lb.) (5 lbs.)

Repacked by
ECONOMY PRODUCTS COMPANY INC.
Shenandoah, Iowa 51601

(back panel)

GENERAL INFORMATION

To mix, slowly sift FORE Turf and Ornamental Fungicide into spray tank ~~and~~ or container while it is being filled, keeping solution well agitated or stirred. Add insecticide and nutritional sprays last. FORE Turf and Ornamental Fungicide is believed compatible with most commonly used pesticides.

DIRECTIONS FOR USE

TURF

Fusarium Blight (F. roseum), Red Thread, Copper Spot, Slime Molds--In a protective spray program use FORE at 4 to 6 oz. per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens up in the spring and repeat every 7 to 14 days.

When conditions are especially favorable for disease development increase FORE Turf and Ornamental Fungicide rate to 6 to 8 oz. per 1000 sq. ft. and repeat applications every 7 days until disease threat is past.

Helminthosporium Melting Out, Rhizoctonia Brown Patch, Rust--Use FORE at 4 oz. per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens-up in the spring and repeat every 7 to 14 days for Helminthosporium and Rust and every 7 days for Rhizoctonia control.

ORNAMENTALS

Use FORE at 1½ level tablespoons per gallon of water (1½ lbs. per 100 gallons) in full coverage spray. Begin spraying when plants are well leafed-out and apply at 7 to 10 day intervals throughout the season for control of:

ROSES - Black Spot; IRIS - Leaf Spot; DAHLIA - Botrytis blight; PANSY - Anthrachnose; ZINNIA - Leaf blight; CARNATION - Leaf spot; ORNAMENTAL CRABAPPLE - Scab, cedar-apple rust; LILY - Botrytis blight; SNAPDRAGON - Rust; TULIP - Botrytis blight (fire); HOLLYHOCK - Leaf Spot, anthrachnose, rust; HOLLY - Purple Spot; HONEYSUCKLE - Herpobasidium blight.

PEONY - Phytophthora and Botrytis blight. Apply in early spring and early fall, drenching soil around plants as well as the foliage. Destroy all infected parts promptly.

FLOWERING DOGWOOD - Anthrachnose. Apply when buds begin to open, when bracts have fallen, 4 weeks later and again in late summer after flower buds for next season are formed.