

DIRECTIONS

To control certain plant disease use this product at rate of 2 tablespoons per gallon of water or 2 lbs. per 100 gallons of water, unless otherwise specified.

APPLES, PEARS — Scab, Frogeye, fruit rots, and blight

— Make application at 5 to 7-day intervals as needed to maintain cover in the pre-bloom, petal-fall, first and second cover sprays. Continue application to within 30 days of harvest to control blight, up to start of harvest on apples to control **fruit rots**. Post-Harvest sprays: Apply immediately after harvest for **leaf spot**. Repeat application in 10 to 14 days. **Fire blight**: Apply at 2 to 3-day intervals during period when new blooms are opening and weather is wet. Maintain recommended sanitary pruning practices. Do not make application to pears within 7 days of harvest.

CHERRY, PEACH, APRICOT, PLUM, GRAPE — On Peach, Plum and Grape apply 2 1/4 lbs. per 100 gallons of water or 2 1/4 tablespoons per gallon of water. On Apricot, use 2 1/2 lbs. per 100 gallons. **Brown rot, cherry leaf spot, peach scab, grape black rot**: Apply in full pink, bloom-petal fall, shuck, cover and post-harvest sprays. Applications at 3 or 4 day intervals may be necessary during bloom to control **blossom blight**. Repeat applications at 7 to 10-day intervals as needed to maintain cover. Do not apply to cherry, grape within 7 days of harvest. Do not apply to peach, plum within 30 days of harvest. Do not apply to apricots between petal fall and harvest. **Post Harvest Sprays**: Apply immediately after harvest for **leaf spot**. Repeat application in 10 to 14 days.

STRAWBERRY — On Strawberries use 2 1/2 lbs. per 100 gallons of water, or 2 1/2 tablespoons per 1 gallon of water. **Botrytis, leaf spot** — Begin when new growth starts in spring. Repeat until fruit begins to form. **Leaf scorch**: Apply in new beds in late summer. Use 2 to 3 weekly applications. Do not apply within 7 days of harvest.

RASPBERRY, BLACKBERRY — Use 2.4 lbs. per acre or 2 1/2 tablespoons per gallon of water. **Anthraco**: Apply thorough coverage in the pre-bloom (whenever new canes are 6 to 8 inches tall) and first cover (right after petal-fall) sprays. Raspberry - **fruit rot**: Begin applications in first bloom. Make 2 repeat applications or more frequently as conditions warrant at approximately 10-15 day intervals. Make thorough coverage, applying spray to point of drip. Make last application 14 days before harvest.

VEGETABLES

Recommended concentrates are 2 level tablespoonfuls per gallon of water or 2 lbs. per 100 gallons of water (unless otherwise specified.) Cover upper and under sides of the leaves thoroughly. Begin application at first sign of infection and repeat at 5 to 7 day intervals or as needed.

BEANS: Rust, anthracnose — Begin when first blossoms appear. Do not apply within 7 days of harvest.

BEETS, CARROTS: Alternaria leaf blight, cercospora leaf spot, downy mildew — Do not apply within 7 days of harvest if crops are to be used as food or feed.

LETTUCE: Downy mildew — apply at weekly intervals. Do not apply within 7 days of harvest.

PEAS: Downy mildew — Begin when first blossoms appear. Do not apply within 10 days of harvest.

CANTALOUPE, CUCUMBERS, WATERMELONS, PUMPKINS, SUMMER and WINTER SQUASH: Downy mildew, Alternaria (Macrosporium) blight, Anthracnose, gummy stem blight.

TOMATO: Early and late blight, anthracnose, stemphylium and septoria leaf spots, Cladosporium gray mold.

PEPPERS: Phytophthora, Anthracnose, Ripe rot.

ORNAMENTALS

LAWNS AND TURF: Brown patch, damping off, leaf spot, root rot, melting out — Use 2 lbs. per 100 gallons or 2 1/2 tablespoons per gallon and apply to surface at the rate of 10 gallons per 1,000 sq. ft. of turf. Repeat at 7 to 10 day intervals as needed. Apply at first signs of disease. For melting out, increase dosage to 3 lbs. per 100 gallons, when grass resumes growth in the spring. Repeat at 7 to 10 day intervals as long as cool, wet weather persists. Do not graze treated areas. Do not feed clippings to livestock.

For the following uses mix 2 level tablespoonfuls per gallon of water. Spray plants thoroughly.

GLADIOLUS: Botrytis blight, curvularia, stemphylium — Spray at weekly intervals or oftener under severe disease conditions, from early stages until spikes begin to open.

ROSES: Black spot — apply at first signs of disease. Repeat at 7 to 10-day intervals.

CARNATIONS, CHRYSANTHEMUMS, SNAPDRAGONS, HOLLYHOCKS: Rust, leaf spot and anthracnose — apply at weekly to 10-day intervals as soon as disease conditions threaten. For control of Botrytis petal spot and Ascochyta-ray make application twice weekly during the blooming season.

AZALEAS and CAMELLIAS: Petal blight — Spray 2 to 3 times a week while flowers are opening. Direct spray into flowers. Spray ground under bushes thoroughly.

PREPLANT DIP FOR SEED POTATOES: seed rot — Mix 2 level tablespoonfuls per gallon of water or 2 lbs. per 100 gallons of water. Immerse pieces in solution for 20-30 minutes. Drain and plant.

PREPLANT DIP FOR GLADIOLUS, TUBEROUS BEGONIAS: seed rot, damping off — Mix 6 to 8 tablespoons per gallon of water. Immerse bulbs in mixture for 20 to 30 minutes. Drain and plant.

NOTICE: Sellers guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 3770-197

SEDA-MBEPCO

MULTI-PURPOSE AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS:

Zineb (zinc ethylene bisdithiocarbamate) 50%

Captan (N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide)* 25%

INERT INGREDIENTS: 25%

*U.S. Patent Nos. 2,553,770; 2,553,771

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid inhalation of dust or spray mist. May cause irritation of nose, throat, and skin. Avoid contact with skin. In case of prolonged or frequent use, wear clean, protective clothing and bathe after each exposure. Do not contaminate food and feed. Keep container closed when not in use. Store in a dry place. Keep away from fire and sparks. Do not store material in any other than original container. Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NET CONTENTS 8 OZ.

Manufactured by:

ECONOMY PRODUCTS COMPANY

SHENANDOAH, IOWA 51601

