

FLY BAIT

With DDVP — 1% VAPONA® INSECTICIDE

**ATTRACTS AND KILLS RESISTANT FLIES,
KILLS MAGGOTS**

ACTIVE INGREDIENTS:		By Weight
2,2-dichlorovinyl dimethyl phosphate		0.93%*
Related Compound		0.07%*
INERT INGREDIENTS:		99.00%
		100.00%


*Equivalent to 1.00% of VAPONA Insecticide RSH-N Trademark

CAUTION

**KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS**

Manufactured By

ECONOMY PRODUCTS CO., INC.
SHENANDOAH, IOWA 51601



GENERAL INFORMATION

FLY BAIT is designed to be used as a supplement to a complete fly control program.

TAKE TIME

OBSERVE LABEL DIRECTIONS

DIRECTIONS

Place baits in areas inaccessible to children and pets.

Dairy Barns, Milk Sheds, Stables, Livestock Barns, Loafing Sheds, Pig Pens and Outdoor Areas: (Flies): Sprinkle **Fly Bait** at the rate of $\frac{1}{4}$ lb. per 1,000 sq. feet on floor or ground area where flies congregate. Ideal baiting sites are around doorways, feed storage rooms, alleyways and window sills. On freshly lined floors place treatment on waste cloth or on paper. Repeat procedure as necessary for practical fly control.

Poultry Houses: (Flies): Sprinkle **Fly Bait** at the rate of $\frac{1}{8}$ lb. (or 2 oz.) per 1,000 sq. ft. where flies congregate but keep out of reach of birds. Apply to matured on window sills, outside of penned enclosures and on floor of feed storage room. Repeat as necessary.

Caged Poultry and Turkeys on Wire: (Flies): Sprinkle **Fly Bait** at the rate of $\frac{1}{8}$ lb. (or 2 oz.) per 1,000 sq. ft. on droppings beneath cages or perches. Repeat at weekly intervals.

Dog Kennels: (Flies): Sprinkle **Fly Bait** at the rate of $\frac{1}{4}$ lb. per 1,000 sq. ft. outside dog runways, on window sills, ledges and other surfaces out of reach of animals. For caged units apply to droppings area. Repeat as necessary.

Garbage Cans: (Flies): Apply **Fly Bait** around and inside garbage can. Repeat as necessary.

For Reducing Populations of Resistant and Non-Resistant Houseflies in and Around Outdoor Living Areas, Picnic Grounds, Backyard Areas, Patios, Loading Docks, Outdoor Latrines, Parking Areas, Refuse Areas Around Service Stations, Open-Air Drive-Ins, Ice Cream Stands, Garbage Collection and Disposal Areas: Use **Fly Bait** at $\frac{1}{2}$ lb. per 1,000 sq. feet of area. Apply on ground or areas where flies congregate. For open-air drive-ins, ice cream stands and restaurants, apply on apron between serving counter and parking area, paying particular attention to trash barrels and garbage cans where flies congregate.

WARNING: Place baits in areas inaccessible to children and pets. **Harmful if Swallowed, Inhaled or Absorbed Through The Skin.** Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Do not reuse empty container. Destroy when empty.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to person or property arising out of use or handling and accepts the product on these conditions.

NET WEIGHT 1 LB.

EPA Reg. No. 3770-202
EPA Est. 3770-1A-1
3770-NB-1

4103B-MBEP00