

Economy

BORDEAUX POWDER

Containing Inert Diluent

CAUTION

KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED

ACTIVE INGREDIENT:

Copper (in Basic Copper Sulfate)

Expressed as Metallic

12.75%

INERT INGREDIENTS

87.25%

NET WEIGHT 1 POUND

Packed by

ECONOMY PRODUCTS CO., INC.

SHENANDOAH, IOWA 51601

DIRECTIONS

Bordeaux Powder should be used in accordance with recommendations of State or Federal Authorities for Basic Copper Sulfate compounds containing the equivalent of 12.75% metallic copper.

FRUIT

Sour Cherries: For Leaf Spot and Brown Rot, use 4 to 5 pounds per 100 gallons of water, or use 3 to 4 tablespoonfuls in one gallon of water. Apply in white bud stage and in late bloom.

Grapes: For Black Rot and Downy Mildew, use 8 pounds per 100 gallons of water, or use 7 tablespoonfuls in one gallon of water. Apply (1) when shoots are 6 to 8 inches long, (2) just after bloom and (3) before the fruit changes color or as local conditions require.

Apples: For late Scab, Blotch and Bitter Rot, use 4 pounds per 100 gallons of water, or use 3 tablespoonfuls in one gallon of water. In areas having severe Blotch or Bitter Rot infections, increase to 8 pounds. Make applications starting at petal fall and continuing through 4th cover or as local conditions require.

CAUTION: Bordeaux Powder is Not recommended for early application on varieties of apples susceptible to copper russetting.

Brambles, Gooseberries and Currants: For Anthracnose and Leaf Spot, use 5½ pounds per 100 gallons of water, or use 4½ tablespoonfuls in 1 gallon of water.

Apply after the leaves open at 7 to 10 day intervals, covering the leaves thoroughly.

Spray as soon as plants become established in the field and repeat applications at 7 to 10 day intervals, as needed, up to harvest.

Potatoes: For Early or Late Blight, use 16 pounds per 100 gallons of water, or 14 tablespoonfuls in one gallon of water.

Tomatoes: For Early and Late Blight and Anthracnose, use 16 pounds per 100 gallons of water, or 14 tablespoonfuls in one gallon of water.

Cucurbits (Cantaloupes, Cucumbers, Squash and Watermelons): For Anthracnose, Downy Mildew, Angular Leaf Spot and Scab, use 8 to 10 pounds per 100 gallons of water, or 7 to 9 tablespoonfuls in 1 gallon of water.

Celery: For Early and Late Blight, use 12 to 16 pounds per 100 gallons of water, or 11 to 14 tablespoonfuls in 1 gallon of water.

NOTICE

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk to persons or property arising out of use or handling and accepts the product on these conditions.

U.S. Reg. No. 100,000

100,000