

POTATO

DUST

For control of Colorado potato beetle, fleabeetles, leafhoppers, blister beetles, cucumber beetles, apply dust lightly but thoroughly to plants once a week.

For control of late blight and early blight on potatoes apply dust lightly but thoroughly to plants once a week when plants are 2-6 inches high. For foliage application only.

IMPORTANT: Keep foliage covered, especially new growth.

CAUTION: May cause irritation of nose, throat, eyes and skin. Do not breathe dust. In case of skin contact wash with water; for eyes get medical attention.

IMPORTANT: Keep away from fire and sparks. Store in a cool, dry place. Do not allow to become wet or overheated in storage. Keep container closed when not in use.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

Empty containers should be disposed of by punching holes in them and burying with wastes or by burning (keep out of smoke).

This guarantee shall be limited to the terms of the label and subject thereto the user assumes all risk to persons or property arising out of use or handling and accepts the product on these conditions.


Feb-26/1970

3776-

213

GURNEY'S

POTATO DUST



ACTIVE INGREDIENTS:

*A coordination product of zinc ion and manganese ethylene bis(dithiocarbamate)	3.9%
In which the ingredients are:	
Manganese ++	7800%
Zinc ++	0975%
Ethylene bis(dithiocarbamate) ion (C ₄ H ₆ N ₂ S ₄)	3.0225%

††Methoxychlor technical 3.0%

INERT INGREDIENTS 93.1%

Total 100.0%

U.S. Pat. Nos. 2,504,404; 2,974,156 and Patent Pending
Equivalent to 2.64% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 0.36% other isomers and reaction products.

KEEP OUT OF REACH OF CHILDREN
See Back Panel for Additional Cautions

NET CONTENTS 1 POUND

Manufactured for

YANKTON, SOUTH DAKOTA 57078