

FLY DEAD SCATTER BAIT

ATTRACTS & KILLS FLIES, KILLS MAGGOTS
WITH VAPONA® AND RONNEL

ACTIVE INGREDIENTS:


2,2-dichlorovinyl dimethyl phosphate	0.228%*
Related Compounds022%*
Ronnel [O,O-dimethyl O-(2,4,5-trichlorophenyl) phosphorothioate]†500%
INERT INGREDIENTS:	99.250%

*Equivalent to 0.25% w of Vapona® Insecticide
†From Technical Grade @Shell Trademark

CAUTION
KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

NET WEIGHT 1 LB.

Manufactured By
ECONOMY PRODUCTS CO., INC.
SHENANDOAH, IOWA 51601



GENERAL INFORMATION

SCATTER BAIT is designed to be used as a supplement to a complete fly control program.

TAKE TIME

OBSERVE LABEL DIRECTIONS

DIRECTIONS

Place baits in areas inaccessible to children and pets.

Caged Poultry and Turkeys on Wire: (Flies): Sprinkle Scatter Bait at the rate of ¼ lb. per 500 sq. ft. on droppings beneath cages or perches. Repeat at weekly intervals.

Poultry Houses: (Flies): Sprinkle Scatter Bait at the rate of ¼ lb. per 500 sq. ft. where flies congregate but keep out of reach of birds. Apply to manure, on window sills, outside of penned enclosures and on floor of feed storage room. Repeat as necessary.

Dog Kennels: (Flies): Sprinkle Scatter Bait at the rate of ¼ lb. per 500 sq. ft. outside dog runways, on window sills, ledges and other surfaces out of reach of animals. For caged units apply to dropping area. Repeat as necessary.

Garbage Cans: (Flies): Apply Scatter Bait around and inside garbage can. Repeat as necessary.

Dairy Barns, Milk Sheds, Stables, Livestock Barns, Loading Sheds, Pig Pens and Outdoor Areas: (Flies): Sprinkle Scatter Bait at the rate of ¼ lb. per 500 sq. feet on floor or ground areas where flies congregate. Ideal baiting sites are around doorways, feed storage rooms, alleyways and window sills. On freshly limed floors place treatment on waste cloth or on paper. Repeat procedure as necessary for practical fly control.

For Reducing Populations of Resistant and Non-Resistant House-flies in and Around Outdoor Living Areas, Picnic Grounds, Back-yard Area, Patios, Loading Docks, Outdoor Latrines, Parking Areas, Refuse Areas Around Service Stations, Open-Air Drive-Ins, Ice Cream Stands, Garbage Collection and Disposal Areas: Use Scatter Bait at ½ lb. per 500 sq. feet of area. Apply on ground or areas where flies congregate. For open-air drive-ins, ice cream stands and restaurants, apply on apron between serving counter and parking area, paying particular attention to trash barrels and garbage cans where flies congregate.

In food processing plants, restaurants, or other places where food is commercially prepared or served, use only when the facility is not in operation and food is not exposed. Use only in bait boxes where bait in each box can be accounted for. Remove and account for all bait boxes prior to resuming food processing.

CAUTION: Place baits in areas inaccessible to children and pets. Harmful if Swallowed, Inhaled or Absorbed Through The Skin. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not reuse empty container. Destroy when empty.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 3770-233
EPA Est. 3770-NB-1

4405B-MBEP CO