

Directions for Use

FACE FLIES: To control face flies, one end of the backrubber must be low enough that it is 15 to 20 inches above ground to allow cattle to rub their heads. The product is only effective for those cattle that rub the face and head. Dilute as for backrubber use.

BACKRUBBER USE: To control hornflies and aid in reducing lice, dilute 1 part Toxaphene with 16 parts kerosene, fuel oil, or motor oil. (Example: 1 cup Toxaphene diluted with 1 gallon oil, 1 gal. Toxaphene diluted with 16 gallons oil). Apply this diluted solution to canvas or burlap cable at the rate of one gallon to 20 feet of cable. Retreat at 2 to 3 week intervals or as necessary to keep cable dampened. Locate backrubber to give beef cattle and hogs free access to it. Best locations are usually near watering places, salt licks or feed racks. **Note:** Do not allow calves under 6 months of age, or dairy animals to treat themselves. Do not permit meat animals access to this treatment within 28 days of slaughter.

BEEF CATTLE, HORSES: Horn Flies, Ticks (Lone Star, Winter and Cattle Fever Ticks) and Lice. Mix 1 gallon Toxaphene with 150 gallons of water. Dip or spray. Spray animals' body thoroughly with enough spray to wet to part the hair and wet the skin. When spraying use about 2 quarts per 100 lb. animal. If reinfestation occurs, repeat application in 14 to 21 days or as needed for control. If flies are the pest, treatment should be repeated until they are completely controlled. Do not treat meat animals for 28 days before slaughter. Do not permit meat animals access to this treatment within 28 days of slaughter.

STOCK TOX

EMULSIFIABLE TOXAPHENE INSECTICIDE

Livestock and Utility Spray

WARNING

KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed. Do not breathe spray mist. May be absorbed through skin. Do not get in eyes, on skin or clothing. Wash thoroughly with soap and water after using. Do not contaminate feed or foodstuffs or drinking water. Do not apply to pets or humans. Do not use on dairy animals. Dip only healthy adult animals under normal conditions. Water animals before dipping. Avoid spraying or dipping animals in cold stormy weather and avoid treating overheated or sick animals. Not for use or storage in or around the home.

Toxic to fish and wildlife; do not contaminate streams, lakes or ponds with this material. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and container. Birds feeding on treated areas may be killed.

Do not reuse empty container. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

ACTIVE INGREDIENTS:

| | |
|---|--------|
| Toxaphene (technical chlorinated camphene*) | 61.20% |
| Xylene | 6.80% |
| Aromatic Petroleum Derivative Solvent | 19.50% |

INERT INGREDIENTS:

| | |
|---------------------------------|---------|
| *Containing 67% to 69% Chlorine | 12.50% |
| | 100.00% |

NET CONTENTS

GALLONS

Manufactured by

ECONOMY PRODUCTS COMPANY, INC.

SHENANDOAH, IOWA 51601

ACCEPTED
10-14-71

3770-246

UNDER THE FEDERAL INSECTICIDE ACT
FOR THE CONTROL OF PESTS
ED
RECOMMENDED COMMENTS

Direction

SHEEP - GOATS: Mix 1 gallon Toxaphene with 150 gallons of water. Dip. When dipping, make sure each animal is completely submerged, making sure that no part of the animal is left out, especially around the neck. Apply during warm weather, before shearing or before replacement ewes are bred before adding to flock. Mix 1 gallon Toxaphene with 150 gallons of water. Dip. If dipping vat is used, agitate before next use.

apply within 28 days of slaughter.

HOGS: Lice. Dilute 1 gallon Toxaphene with 150 gallons of water. Dip within 30 days of farrowing. Spraying prevents pig lice. Do not treat pigs until 28 days before slaughter. Do not treat meat hogs.

SARCOPTIC MANGE: Dilute 1 gallon Toxaphene with 150 gallons of water. Dip sheep and goats for 14 days. Apply to head and neck. **Note:** Do not use on small animals and do not treat meat animals before slaughter.

UTILITY SPRAY: To control the pest, spray with 1 gallon Toxaphene with 150 gallons of water. Spray utility animals, and spray utility animals. Do not treat meat animals before slaughter.

NOTICE: Do not use on small animals and do not treat meat animals before slaughter.

STOCK TOX

EMULSIFIABLE TOXAPHENE INSECTICIDE

Livestock and Utility Spray

ACCEPTED
10-14-71
3770-246
UNDER THE FEDERAL INSECTICIDE ACT
FOR THE CONTROL OF PESTS

WARNING

KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed. Do not breathe spray mist. May be absorbed through skin. Do not get in eyes, on skin or clothing. Wash thoroughly with soap and water after using. Do not contaminate feed or foodstuffs or drinking water. Do not apply to pets or humans. Do not use on dairy animals. Dip only healthy adult animals under normal conditions. Water animals before dipping. Avoid spraying or dipping animals in cold stormy weather and avoid treating overheated or sick animals. Not for use or storage in or around the home.

Toxic to fish and wildlife; do not contaminate streams, lakes or ponds with this material. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and container. Birds feeding on treated areas may be killed.

Do not reuse empty container. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

ACTIVE INGREDIENTS:

| | |
|---|--------|
| Toxaphene (technical chlorinated camphene*) | 61.20% |
| Xylene | 6.80% |
| Aromatic Petroleum Derivative Solvent | 19.50% |

INERT INGREDIENTS:

| | |
|---------------------------------|---------|
| *Containing 67% to 69% Chlorine | 100.00% |
|---------------------------------|---------|

NET CONTENTS

GALONS

Manufactured by
ECONOMY PRODUCTS COMPANY, INC.
SHENANDOAH, IOWA 51601

Directions for Use

SHEEP and GOATS: Lice: Dilute 1 gallon Toxaphene in 350 gallons of water. Dip or spray thoroughly. (When dipping make sure the animal is held in vat at least 2 minutes and submerge each animal twice during dipping, making sure that material penetrates wool specially around the head. For **Sheep Ticks** apply during warm weather soon after shearing or before wool is 2 inches long. Replacement ewes or rams should be treated before adding to flock. For **Sheep Scabies** mix 1 gallon Toxaphene to 150 gallons water. If dipping vat is intermittently used, agitate before next dipping. Do not dip or spray within 28 days of slaughter. Do not apply to dairy goats. Do not use on animals under 3 months of age.

HOGS: Lice: Dilute 1 gallon Toxaphene in 350 gallons of water. Do not treat sows within 30 days of farrowing nor for three weeks thereafter. Spraying sows before farrowing prevents pigs from being infested. Do not treat pigs under 6 months of age. Do not treat meat hogs within 28 days of slaughter.

SARCOPTIC MANGE: Beef cattle, sheep and goats: Mix 1 gallon Toxaphene in 150 gallons of water. Dip or spray cattle, dip sheep and goats, for thorough animal coverage. Apply to herd when mange is found. Repeat treatment in 15-21 days if necessary. Note: Do not apply to dairy animals or small animals under 6 months of age. Do not treat meat animals within 28 days of slaughter.

UTILITY SPRAY: For mosquitoes, flies, gnats. To control these pests in outbuildings, sheds, barns, dilute 1 gallon Toxaphene with 11 gallons of water. Make application to walls, door frames, ground areas, ceiling and places where insects rest or breed. Do not use on dairy farms, milk horses or poultry houses. If spot application is needed for control of flies, the spray should be applied to the animal's head, neck, and body. Do not use on animals under 3 months of age. Do not use on animals under 28 days of slaughter.

NOTICE: Some states have laws which require the use of certain insecticides. It is the user's responsibility to determine the laws of his state and to use the insecticide in accordance with the law.