DIRECTIONS

VAPONA Insecticide is a very effective fog or surface spray and will provide fast knockdown and kill of flies, mosquite is and other small flying insects. It is a multi-purpose formulation designed for use as an animal spray to protect dairy cows, beef cattle and horses from face flies, stable flies, house flies, horn flies, gnats and mosquitoes.

USE AS A BARN FOG - VAPONA Insecticide can be used in barn fogging equipment without dilution. One pint will fog 8,000 cubic feet of enclosure. Before applying, reduce air movement as much as possible by closing doors, windows and other openings. Do not contaminate water, feed or foodstuffs, milk or milking utensils. Do not use this solution for fogging purposes in areas where animals have received direct application within 8 hours.

ANIMAL BUILDINGS (Horse Barns, Dairy Barns, Shelter Sheds), MILK SHEDS, OUTDOOR AREAS ___ Dilute VAPONA Insecticide with equal amounts of deodorized kerosene or fuel oil. Apply as a coarse wet spray at the rate of one pint per 1,000 square feet to walls, ceilings, stalls, feed storage rooms, alleyways, pig pens, milk sheds, loafing areas. In poultry houses apply to manure, window sills and outside of penned enclosures. For fly and maggot control in breeding areas such as dumps and manure piles apply at the rate of 1 to 2 quarts of the diluted mixture per 100 sq. ft. Apply this diluted mixture as a coarse wet spray at the rate of one quart per 1.000 sq. feet to treat loading docks, parking areas, refuse areas around service stations, open air drive-ins, garbage collection and disposal areas.

VAPONA® INSECTICIDE

31110 -160 READY TO USE - CONTAINS 1% VAPONA (DDVP) Related Compounds 100. Equivalent to 1.00% w of Vapona Insecticide RShell Chemical Co THE PROFESAL INCRETICIOE FOR LACTATING DAIRY CATTLE, BEEF CATTLE AND BA FUNDICIDE AND ROBENTICIDE ACT

FOR ECONOMIC POISON REGISTER-

TO ATTACHED COMMENTS.

SHONECE

ED UNDER NO.

WARNING

KEEP OUT OF REACH OF CHILDREN

May Be Fatal If Swallowed, Inhaled Or Absorbed Through The Skin. Do not or on skin. Wash thoroughly with soap and water after handling and before smoking. In case of contact with skin or eyes, flush with plenty of water, medical attention. Do not contaminate feed, water, milk or milk handling (swallowed, induce vomiting immediately and get medical attention. Atropine

This product is toxic to fish and wildlife. Birds feeding on treated areas m Keep out of any body of water. Apply this product only as specified on the Do not reuse empty container. Destroy it by perforating or crushing. Bur in a safe place.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN

NET CONTENTS 1 GALLON

Manufactured By ECONOMY PRODUCTS COMPANY, INC SHENANDOAH, IOWA 51801

VAPONA® INSECTICIDE

LEADY TO USE -- CONTAINS TO VAPONA (DOVP)

E INGREDIENTS: lichlorovinyl dimethyl phosphate led Compounds leum Oil	0.07%*
	100.00%

alent to 1.00% w of Vapona Insecticide

R Shell Chemical Company

LACTATING DAIRY CATTLE, BEEF CATTLE AND BARNS

KEEP OUT OF REACH OF CHILDREN

If Swallowed, Inhaled Or Absorbed Through The Skin. Do not get in eyes Vash thoroughly with soap and water after handling and before eating or ase of contact with skin or eyes, flush with plenty of water, for eyes get tion. Do not contaminate feed, water, milk or milk handling equipment. If duce vomiting immediately and get medical attention. Atropine is antidotal.

is toxic to fish and wildlife. Birds feeding on treated areas may be killed. iny body of water. Apply this product only as specified on this label.

empty container. Destroy it by perforating or crushing. Bury or discard

USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

NET CONTENTS 1 GALLON

Manufactured By

SHENANDOAH, IOWA \$1601

DIRECTIONS

Apply VAPONA Insecticide as a mist spray with hand or automatic sprayer for control of face flies, stable flies, house flies, horn flies, gnats and mosquitoes.

USE AS AN ANIMAL SPRAY --- On dairy cows, beer cattle and horses, apply 2 fluid ounces of VAPONA Insecticide to each animal each day. The mist spray should be applied with a good hand or automatic sprayer. Spray to wet hair. For maximum results apply the 2 fluid ounces in 2 applications-1 ounce morning and night. Special care should be used to assure coverage of the legs, belly, shoulders and head. Caution: Do not apply regularly to calves under 6 months of age. Apply thoroughly as a mist spray to animals but do not wet skin. Keep unprotected persons out of operating area. In a complete fly control program this spray should be used in conjunction with a VAPONA Insecticide building, fog or surface spray and sanitary measures. I'm not use this solution for fogging purposes within 8 hours of direct spray to animals.

NOTICE — Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 3770-260

±116B-MBEPCO