

DIRECTIONS FOR USE

To develop a residual repellency to stable flies, horn flies, mosquitoes and gnats on dairy cows, apply 1 to 2 ounces of **Dairy Cattle Spray** to each animal for a three day period. The spray should be applied with a good hydraulic-type hand sprayer so that the animal's entire body has been uniformly covered. Special care should be used in spraying the legs, belly and shoulders. Avoid areas around the nose, eyes and mouth. Avoid wetting the hide. Spray in the same manner again after two days have elapsed and every second day thereafter or when insects again disturb the animal. Be sure the teats and udder are washed with warm water prior to milking. Calves should be sprayed very lightly once a day instead of attempting to build up a residual repellency by spraying 2 or more heavily.

For control of lice on livestock, spray the animals with 2 ounces of **Dairy Cattle Spray**. Immediately rub the spray into the hair, paying particular attention to infested areas. Repeat as necessary, but not more frequently than once every three weeks.

To kill flies, mosquitoes and gnats in the barn prior to milking, close all doors and windows and direct the spray toward the ceiling. The barn should remain closed for 10 to 20 minutes. Sweep up and destroy fallen insects. Repeat as necessary. Apply at the rate of one ounce spray per 1,000 cubic feet. Do not use this mist solution in areas where animals have received direct spray within 8 hours. Do not contaminate water, feed or foodstuffs, milk or milking utensils.

To protect livestock from attack by fleas, spray bedding and walls of pen, particularly cracks and crevices. Repeat treatment frequently until adequate control is obtained.

For complete flea control on dogs, this spray should be used in conjunction with residual sprays on heads and on their fleas.

NOTICE

See the label for full directions for use. Do not use this product on any animal unless the product is specifically labeled for that use.

TRADE MARK

10
E
W
S
T
E
R
N
I
N
D
U
S
T
R
I
A
L
C
O
M
P
A
N
Y

DAIRY CATTLE SPRAY

Kills and repels horn flies, stable flies, mosquitoes and gnats. When properly applied, the repellent effect against above insects will last two days.

ACTIVE INGREDIENTS:	100%
Pyrethrins	.05%
*Piperonyl butoxide, technical	.10%
**N-octyl bicycloheptene dicarboximide	.16%
Di-n-propyl isocinchomeronate	.20%
2,2-Dichlorovinyl dimethyl phosphate	.465%
Related Compounds	.035%
Petroleum distillate	98.990%
*Consists of 0.08% (butylcarbityl) (6-propylpiperonyl) ether and 0.02% other related compounds	
**MGK 264, insecticide Synergist, U. S. Pat. No. 2,476,512	
MGK Repellent 326, U. S. Pat. No. 2,757,120	
Equivalent to .50% w of Vapona® Insecticide - Shell Trademark	

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid prolonged breathing of spray mist.

Harmful if swallowed. Consult a physician immediately.

Store in a cool place. Keep container tightly closed when not in use. Avoid contamination or milk and milk handling equipment. Avoid overspraying which will result in wetting the animals' hides. Do not reuse empty container. Destroy when empty.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

NET CONTENTS 1 GALLON

Manufactured by:

ECONOMY PRODUCTS COMPANY, INC.

SHELANDOAH, IOWA 51601

DIRECTIONS FOR USE

To develop a residual repellency to stable flies, horn flies, mosquitoes and gnats on dairy cows, apply 1 to 2 ounces of **Dairy Cattle Spray** to each animal for a three day period. The spray should be applied with a good hydraulic-type hand sprayer so that the animal's entire body has been uniformly covered. Special care should be used in spraying the legs, belly and shoulders. Avoid areas around the nose, eyes and mouth. Avoid wetting the hide. Spray in the same manner again after two days have elapsed and every second day thereafter or when insects again disturb the animal. Be sure the teats and udder are washed with warm water prior to milking. Calves should be sprayed very lightly once a day instead of attempting to build up a residual repellency by spraying more heavily.

For control of lice on livestock, spray the animals with 2 ounces of **Dairy Cattle Spray**. Immediately rub the spray into the hair, paying particular attention to infested areas. Repeat as necessary, but not more frequently than once every three weeks.

To kill flies, mosquitoes and gnats in the barn prior to milking, close all doors and windows and direct the spray toward the ceiling. The barn should remain closed for 10 to 20 minutes. Sweep up and destroy fallen insects. Repeat as necessary. Apply at the rate of one ounce spray per 1,000 cubic feet. Do not use this mist solution in areas where animals have received direct spray within 8 hours. Do not contaminate water, feed or foodstuffs, milk or milking utensils.

To protect livestock from attack by fleas, spray bedding and walls of pen, particularly cracks and crevices. Repeat treatment frequently until adequate control is obtained.

For complete flea control on dogs, this spray should be used in conjunction with residual sprays on heads and on their fleas.

NOTICE

See the label for full directions for use. Do not use this product on any animal unless the product is specifically labeled for that use.

TRADE MARK