

DIRECTIONS (Vegetables, Small Fruit, Ornamentals)

Thorough, full-coverage applications should be made. Dust underneath and top of foliage thoroughly. Apply to both plants and insects. Begin application when plants are well established or when new growth appears. Repeat application as necessary.

CONTROL CHART

Apply before plants are infested and damaged by the following insects:

7-Day Interval Between Last Application and Harvest:

Cabbage, Kale, Brussels Sprouts, Mustard Greens

(Aphids, Imported Cabbage Worm, Cabbage Looper):

Carrots, Head Lettuce (Aphids, Leafhoppers): **Caull-**

flower (Aphids): **Radishes, Parsnips, Salsify, Horse-**

radish, Swiss Chard, Spinach (Aphids): **Peas** (Pea

Aphids) Apply to foliage during early bloom period before eggs are laid; repeat as necessary.

5-Day Interval Between Last Application and Harvest:

Sweet Corn (Aphids).

3-Day Interval Between Last Application and Harvest:

Broccoli, Turnip (Aphids, Imported Cabbage Worm, Cabbage

Looper): **Onions** (Thrips): **Pumpkins** (Squash Vine

Borer) Apply once or twice a week to stems and vines

at base of plants. Do not apply to pumpkins unless

plants are dry: **Strawberries** (Lygus Bug, Spittlebugs,

Field Crickets, Thrips, Potato Leafhopper, Strawberry

Leafhopper): **Peppers** (Pepper Maggots): **Sweet Potato**

(Leafhopper).

1-Day Interval Between Last Application and Harvest:

Asparagus (Thrips): **Beans** (Mexican Bean Beetle)

Cucumbers, Melons (Aphids, Leafhopper, Spider Mite,

Cucumber Beetle) Do not apply to cucumbers, melons

unless plants are dry: **Squash** (Aphids, Spider Mites,

Squash Vine Borer, Cucumber Beetle) Apply weekly to

stems and vines at base of plants. Do not apply to

squash unless plants are dry: **Tomato** (Spider Mites,

Aphids, Tomato Russet Mite, Drosophila): **Blackberries,**

Boysenberries (Aphids).

No time limitation: **Potatoes** (Aphids, Leafhopper,

Mealybug). For Mealybug, apply on foliage of infested

plants and repeat as necessary.

Strawberry (Strawberry Root Weevil) Apply to soil

surface before planting and work into top 6 to 8 inches

of soil. Apply at the rate of 10-15 ounces per 1,000

sq. feet of soil area (equivalent to 27 to 40 lbs. per

acre).

Ornamentals (Aphids, Spider Mites) Make thorough,

full-coverage applications. Repeat applications as neces-

sary. Injury may occur on certain ferns such as Boston,

Maidenhair and on some species of crassula and

canaerti Juniper.

4% MALATHION POWDER INSECTICIDE

ACTIVE INGREDIENT:

Malathion* 4%

INERT INGREDIENTS 96%

*O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate

CAUTION KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed or inhaled. Avoid breathing of dust. Avoid contact with skin. Wash thoroughly with soap and water after handling. Avoid contamination of feed and foodstuffs. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

ENVIRONMENTAL CAUTION: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

CONTAINER DISPOSAL: Do not reuse container. Destroy when empty.

NET WEIGHT 1 LB.

Manufactured by:

ECONOMY PRODUCTS CO., INC.

SHENANDOAH, IOWA 51601

ACCEPTED

APR 23 1975

Under the Federal Insect
Fungicide,
as a

DIRECTIONS

(Animals)

3770-277

POULTRY: (Northern Fowl Mite, Poultry Lice) Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4 to 8 weeks or when necessary.

HOGS: (Lice) Treat animals thoroughly. Repeat in 10 days and thereafter as needed.

DOGS and CATS: (Fleas) Direct application. Apply sufficient dust to treat animal thoroughly. Use any garden-type duster, puff duster or shaker-type can. If animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animals should be closely examined for fleas between treatment. If the animal is confined to an area free of fleas, one treatment may be sufficient.

CATTLE (Beef and Non-Milking): (Lice) Treat animal thoroughly. Repeat in 10 days and thereafter as needed. (Horn Fly) Use 4 tbs. per animal. Apply on back and neck of the animal and repeat applications at 10-14 day intervals. Note: Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening.

DAIRY CATTLE: (Horn Fly) Use 4 tbs. per animal. Apply on back and neck of the animal. Do not apply to dairy cattle later than 5 hours before milking or during milking. Repeat applications at 10- to 14-day intervals if needed.

Do not treat animals under one month of age. When applying dusts avoid contamination of feed, food containers and watering troughs.

NOTICE TO USERS: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 3770-277

EPA Est. 3770-NB-1

3985B-MBEPKO