



Camerman's Note
Poor Copy

GENERAL INFORMATION — Apply with hand power or automatic equipment as a mist spray to moisten the hair only and not in excess of 2 fluid ounces daily per animal. Do not wet the skin due to possible skin irritation. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates. Keep unprotected persons out of operating area.

In a complete fly control program this spray should be used in conjunction with other sanitary measures.

DIRECTIONS

Lactating Dairy Cows, Beef Cattle,
Horses, Hogs

FACE FLIES: To protect animals from Face Flies, spray animal's head, covering forehead, poll, muzzle and sides of face thoroughly with fine mist but do not apply more than 1 oz. to entire animal at any one spraying.

STABLE FLIES, HORN FLIES, HOUSE FLIES, MOSQUITOES: For fast knock-down and kill of pests, apply as a mist over milk cows, horses, beef cattle, hogs. Kills stable flies, horn flies, house flies, mosquitoes, black flies and sand flies at time of spraying. For maximum results apply 2 fluid ounces in 2 applications — 1 ounce morning and evening.

(Continued on other side)

LIVESTOCK - DAIRY SPRAY AND FOG

With Vapona (DDVP) and Ciodrin
Ready-to-use

KILLS

Face Flies, Stable Flies, House Flies,
Horn Flies, Fleas and Mosquitoes

USE ON

Lactating Dairy Cows, Beef
Cattle, Horses, Hogs

ACTIVE INGREDIENTS:

2,2-Dichlorovinyl dimethyl phosphate	0.02%	.93
Related compounds	0.08%	.07
Dimethyl phosphate of alpha methylbenzyl-3-hydroxy-cis-crotonate**	0.25%	.25
Petroleum distillate	49.00%	49.69
Mineral oil	49.00%	49.00
INERT INGREDIENTS:	0.00%	0.00

*Equivalent to 1.00% VAPONA Insecticide
**CIODRIN Insecticide
© Shell Chemical Company

WARNING

KEEP OUT OF REACH OF CHILDREN

May Be Fatal If Swallowed, Inhaled Or Absorbed Through The Skin. Do not get in eyes or on skin. Wash thoroughly with soap and water after handling and before eating or smoking. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Do not contaminate feed, water, milk or milk handling equipment.

If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

NET CONTENTS 1 GALLON

Manufactured by

ECONOMY PRODUCTS COMPANY, INC.

SHELANDOAH, IOWA 51601

Special care should be used to assure complete coverage of the legs, body, shoulders and head. For best results, spray milk cows before daily milking time for 7-10 days. Apply 1 oz. per head at morning and evening milkings. Complete coverage is necessary for best possible protection. Treatments can be discontinued until flies reappear on animals.

ELECTRIC and STATIONARY FOGGERS: Livestock-Dairy Spray and Fog is recommended for use in stationary or electric foggers. Fog about 2 ounces per 1,000 cubic feet of enclosure. Before using fog can be applied around each animal. Before applying, reduce air flow as much as possible by closing doors, windows and other openings. Then treated area should be closed for 10 minutes. Do not apply in areas where animals have received daily applications of DDVP within 8 hours.

FLEAS: Spray bedding and sleeping quarters where infested animals are housed. Spray thoroughly floors, walls, cracks, crevices, sleeping quarters of animals or other infested areas. Repeat application if necessary.

NOTICE: Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons or property arising out of use or misuse and accepts the product on these terms.

EPA Reg. No. 3770-282

51601 A