

Potato Dust  
DIRECTIONS

POTATOES (early and late blights): Apply 11-2/3 to 20 lbs. (1 3/4 to 3 lbs. actual Captan) as a dust per acre. Begin when plants are 2 to 4 inches high. Repeat at 5- to 7-day intervals throughout the season. No time limitation between last application and harvest.

Protect from excessive heat  
Store in Cool, Dry Place

EPA Reg. No. 3770-285

EPA Est. 3770-1A-1



3770-NB-1 ←

15% CAPTAN DUST  
For Disease Control  
on Potatoes

ACTIVE INGREDIENT:  
Captan: (N-trichloromethylthio-  
4-cyclohexene-1,2-dicarboxi-  
mide) 15%  
INERT INGREDIENTS: 85%

CAUTION  
KEEP OUT OF REACH  
OF CHILDREN

Avoid inhaling dust. Wear respirator.  
Avoid contact with skin. Wash with  
soap and water after handling. Do not  
contaminate feed, foodstuffs, or water.  
See side panel for additional cautions.

Net Weight \_\_\_\_\_

Manufactured by:  
**Economy Products  
Company Inc.**

Shenandoah, Iowa 51601

ENVIRONMENTAL CAUTION: This product is toxic to fish, birds, and other wildlife. Keep out of any lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from the areas treated.

CONTAINER DISPOSAL: Do not reuse container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTICE TO USER: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

ACCEPTED

*Detol 24, 1974*

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 3770-285