


E P C O

TRICHLORFON 80% SOLUBLE POWDER

ANIMAL INSECTICIDE

FOR CONTROL OF HORN FLIES, LICE AND GRUBS ON BEEF CATTLE

DIRECTIO



GENERAL INFORMATION—

Trichlorfon 80% Soluble Powder is a highly effective insecticide that can be used in any type of conventional spray equipment. Trichlorfon 80% Soluble Powder completely dissolves in water for a more effective spray solution. Trichlorfon 80% Soluble Powder gives systemic control of grubs and gives residual control of lice and horn flies. Repeat treatments should not be made more often than at 7-day intervals.

For Cattle Grub Control — Treatment of cattle with Trichlorfon 80% Soluble Powder can be made any time during the grub season. However, for best results, treat early in the grub cycle. Treat as soon as possible after activity of heel fly has stopped, but at least 6 weeks prior to expected appearance of grubs. You may apply Trichlorfon 80% Soluble Powder very near or during the time grubs appear, but considerable grub damage will have already occurred. Use as spray treatment. Mix 10 lbs. of Trichlorfon 80% Soluble Powder in 100 gallons of water. Apply this solution for complete wetting and run-off. Apply in a manner that the skin, not the hair, becomes thoroughly wet. Spray pressure of 250 to 350 lbs. are suggested. Operate box-type spray chutes at maximum pressure for thorough over-all wetting and penetration. A single application is usually adequate for grub control. Do not use within 14 days of slaughter. Consult State Agricultural Extension Service personnel for additional information concerning proper timing of applications for cattle grub control.

Cattle Horn Fly Control — Use when horn flies appear and constitute a problem. Spray treatment. Mix 10 lbs. of Trichlorfon 80% Soluble Powder in 100 gallons of water. Apply this solution for complete wetting and run-off. Repeat as necessary, but do not use within 14 days of slaughter.

Cattle Lice Control — Use when lice appear and constitute a problem. Use as spray treatment: 2½ lbs. of Trichlorfon 80% Soluble Powder in 100 gallons of water. Apply this solution for complete wetting and run-off. Repeat as necessary but do not use within 14 days of slaughter.

Do not treat dairy animals. Do not treat animals less than 3 months old. Do not treat animals that are sick, convalescent, or stressed livestock. Do not spray in a confined, non-ventilated area. Do not use in conjunction with oral drenches, other medications, or other organic phosphate-type sprays.

ACTIVE INGREDIENT:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate* .. 80%

INERT INGREDIENTS: 20%

Total 100%

*Trichlorfon

WARNING

KEEP OUT OF REACH OF CHILDREN


See Back Panel for Additional Warnings

NET WEIGHT ———

Manufactured For:

ECONOMY PRODUCTS COMPANY, INC.
SHENANDOAH, IOWA 51601

DIRECTIONS FOR USE



GENERAL INFORMATION—

Trichlorfon 80% Soluble Powder is a highly effective insecticide that can be used in any type of conventional spray equipment. Trichlorfon 80% Soluble Powder completely dissolves in water for a more effective spray solution. Trichlorfon 80% Soluble Powder gives systemic control of grubs and gives residual control of lice and horn flies. Repeat treatments should not be made more often than at 14-day intervals.

For Cattle Grub Control — Treatment of cattle with Trichlorfon 80% Soluble Powder can be made any time during the grub season. However, for best results, treat early in the grub cycle. Treat as soon as possible after activity of heel fly has stopped and at least 6 weeks prior to expected appearance of grubs. You may apply Trichlorfon 80% Soluble Powder very near or during the time grubs appear but considerable grub damage will have already occurred. Use as spray treatment. Mix 10 lbs. of Trichlorfon 80% Soluble Powder in 100 gallons of water. Apply this solution for complete wetting to run-off. Apply in a manner that the skin, not just the hair, becomes thoroughly wet. Spray pressures of 250 to 350 lbs. are suggested. Operate box-type spray chutes at maximum pressure for thorough over-all wetting and penetration. A single application is usually adequate for grub control. Do not use within 14 days of slaughter. Consult State Agricultural Extension Service personnel for additional information concerning proper timing of applications for cattle grub control.

Cattle Horn Fly Control — Use when horn flies appear and constitute a problem. Spray treatment: Mix 10 lbs. of Trichlorfon 80% Soluble Powder in 100 gallons of water. Apply this solution for complete wetting and run-off. Repeat as necessary but do not use within 14 days of slaughter.

Cattle Lice Control — Use when lice appear and constitute a problem. Use as spray treatment: mix 2½ lbs. of Trichlorfon 80% Soluble Powder in 100 gallons of water. Apply this solution for complete wetting to run-off. Repeat as necessary but do not use within 14 days of slaughter.

Do not treat dairy animals. Do not treat animals less than 3 months old. Do not treat animals that are sick, convalescent, or stressed livestock. Do not spray in a confined, non-ventilated area. Do not use in conjunction with oral drenches, other medications, or other organic phosphate-type sprays.

WARNING:

Trichlorfon is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

Atropine is antidotal.

Consult veterinarian at first sign of an adverse reaction.

Do not reuse empty container. Destroy empty container by perforating or crushing and burying in a safe place.

WARNING

Keep away from domestic animals and foodstuffs.

This product is toxic to fish, wildlife and birds. Do not contaminate water by cleaning of equipment or disposal of wastes.

May be harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust or spray mist. Always wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and warm water before re-use. Do not contaminate feed or food.

If poisoning occurs, get medical attention immediately. Prolonged exposure will result in cholinesterase depression.

Note to Physician: Atropine sulfate is antidote. 2-PAM is also antidote and may be used in conjunction with atropine.

NOTICE

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.