

DIRECTIONS FOR LIVESTOCK PEST CONTROL

For Control of Face Flies, Horn Flies, House Flies, Stable Flies, Lice, and Lone Star Ticks on Dairy Cattle, Beef Cattle, Sheep, Goats and Hogs. Mix 2 quarts of CI-PONA Insecticide Emulsifiable per 6 gallons of water. Spray to thoroughly cover all parts of the animal, including the legs, using 1-2 pints of spray per large animal and proportionately less for smaller animals.

For pen or corral operations involving spray application to livestock with high pressure equipment, do not exceed a 0.5% concentration. Mix 2 quarts of CI-PONA Insecticide Emulsifiable per 12 gallons of water (0.5% concentration). Spray to thoroughly cover the animal's body and legs (avoid excessive spraying of the head) using 1-2 quarts of spray per large animal and proportionately less for smaller animals. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

Improved face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gallons of spray. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

Low Volume Application for Control of Face Flies. Mix 4 tablespoons (2 fluid oz.) of CI-PONA Insecticide Emulsifiable per 1½ pints of water (1% concentration). Using a calibrated sprayer, apply 5 teaspoons of spray to the face and a total of 6-10 teaspoons to the back and sides of the animal 3 times per week. Improved control may be achieved by the addition of 1 level teaspoon of sugar per pint of spray solution.

For Control of Lice and Winter Ticks on Dairy Cattle, Beef Cattle, Sheep, Goats and Hogs. Mix 2½ pints of CI-PONA Insecticide Emulsifiable per 16 gallons of water (0.25% concentration). Spray animals thoroughly using up to 1 gallon of spray. Apply a second application 14 days later. Repeat applications as necessary to maintain control, but not more often than once every 7 days.

Backrubber Applications. For use in backrubbers or face rubbers for the control of horn flies and face flies, mix 1 quart of CI-PONA Insecticide Emulsifiable per 4 gallons of back rubber base oil (1% concentration). This may be diesel fuel or fuel oil in accordance with local recommendations.

For Control of House Flies, Face Flies, and Stable Flies in and Around Livestock Buildings. Mix 2 quarts of CI-PONA Insecticide Emulsifiable per 6 gallons of water (1% concentration). Spray premises such as inside and outside barn walls (including dairy barns), eaves, fences and other resting or breeding places at least twice monthly using approximately 1 gallon of finished spray per 1,000 square feet (40'x25'). Improved house fly and

(Continued on other side)

CI-PONA Dairy & Livestock Insecticide

EMULSIFIABLE LIQUID

CONTAINS CIODRIN® AND VAPONA® INSECTICIDES

Dilute With Water or Oil

TAKE TIME



OBSERVE LABEL DIRECTIONS

ACTIVE INGREDIENTS:	By Weight
Dimethyl phosphate of alpha-methylbenzyl	10.0%
3-hydroxy-cis-crotonate	2.3%
2,2-dichlorovinyl dimethyl phosphate	9.2%
Related Compounds	77.0%
Petroleum Hydrocarbons	10.5%
INERT INGREDIENTS:	10.5%
Total	100.0%
*Equivalent 2.5% VAPONA® Insecticide	
©Shell Chemical Company	

WARNING

KEEP OUT OF REACH OF CHILDREN
See Side Panel for Additional Cautions

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

NET CONTENTS _____

Manufactured by

ECONOMY PRODUCTS COMPANY, INC.

SHEMANDOAN, IOWA 51601

DIRECTIONS FOR LIVESTOCK PEST CONTROL (Continued)

face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gallons of spray.

Do not use in poultry houses.

For Control of Chorioptic Mites on Dairy and Beef Cattle. Mix 2½ pints of CI-PONA Insecticide Emulsifiable per 16 gallons of water (0.25% concentration). Spray animals thoroughly. Retreat within 10-14 days for lasting control.

WARNING

WARNING! HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

During commercial or prolonged exposure in spray-misting and loading operations, wear clean rubber gloves and a mask or respirator of a type found suitable by the U. S. Bureau of Mines for protection against CIODRIN-VAPONA insecticide. Use this material only for recommended purposes and at recommended dosages.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not treat animals within one day of slaughter.


Do not apply dermally in combination with Trichlorfon.

DECONTAMINATE AND DESTROY USED CONTAINERS. Do not re-use for any purpose. Decontaminate containers by rinsing thoroughly with water and alkaline detergent. Destroy containers by breaking and burying fragments in isolated area. Dispose of rinsings in a way as not to constitute a hazard, or contaminate water supplies.

NOTICE: Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 8770-802
EPA Est. 8770-1A-1
8770-NB-1

2919B-MBEPCCO



DIRECTIONS FOR LIVESTOCK PEST CONTROL

For Control of Face Flies, Horn Flies, House Flies, Stable Flies, Lice, and Lone Star Ticks on Dairy Cattle, Beef Cattle, Sheep, Goats and Hogs. Mix 2 quarts of CI-PONA Insecticide Emulsifiable per 6 gallons of water. Spray to thoroughly cover all parts of the animal, including the legs, using 1-2 pints of spray per large animal and proportionately less for smaller animals.

For pen or corral operations involving spray application to livestock with high pressure equipment, do not exceed a 0.5% concentration. Mix 2 quarts of CI-PONA Insecticide Emulsifiable per 12 gallons of water (0.5% concentration). Spray to thoroughly cover the animal's body and legs (avoid excessive spraying of the head) using 1-2 quarts of spray per large animal and proportionately less for smaller animals. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

Improved face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gallons of spray. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

Low Volume Application for Control of Face Flies. Mix 4 tablespoons (2 fluid oz.) of CI-PONA Insecticide Emulsifiable per 1½ pints of water (1% concentration). Using a calibrated sprayer, apply 5 teaspoons of spray to the face and a total of 5-10 teaspoons to the back and sides of the animal 3 times per week. Improved control may be achieved by the addition of 1 level teaspoon of sugar per pint of spray solution.

For Control of Lice and Winter Ticks on Dairy Cattle, Beef Cattle, Sheep, Goats and Hogs. Mix 2½ pints of CI-PONA Insecticide Emulsifiable per 16 gallons of water (0.25% concentration). Spray animals thoroughly using up to 1 gallon of spray. Apply a second application 14 days later. Repeat applications as necessary to maintain control, but not more often than once every 7 days.

Backrubber Applications. For use in backrubbers or face rubbers for the control of horn flies and face flies, mix 1 quart of CI-PONA Insecticide Emulsifiable per 4 gallons of back rubber base oil (1% concentration). This may be diesel fuel or fuel oil in accordance with local recommendations.

For Control of House Flies, Face Flies, and Stable Flies in and Around Livestock Buildings. Mix 2 quarts of CI-PONA Insecticide Emulsifiable per 6 gallons of water (1% concentration). Spray premises such as inside and outside barn walls (including dairy barns), eaves, fences and other resting or breeding places at least twice monthly using approximately 1 gallon of finished spray per 1,000 square feet (40'x25'). Improved house fly and

(Continued on other side)

CI-PONA Dairy & Livestock Insecticide

EMULSIFIABLE LIQUID

CONTAINS CIODRIN® AND VAPONA® INSECTICIDES

Dilute With Water or Oil

TAKE TIME



OBSERVE LABEL DIRECTIONS

ACTIVE INGREDIENTS:	By Weight
Dimethyl phosphate of alpha-methylbenzyl	10.0%
3-hydroxy-cis-crotonate	2.5%
2,2-dichlorovinyl dimethyl phosphate	2.5%
Related Compounds	0.2%
Petroleum Hydrocarbons	77.8%
INERT INGREDIENTS:	10.5%
Total	100.0%
*Equivalent 2.5% VAPONA® Insecticide	
©Shell Chemical Company	

WARNING

KEEP OUT OF REACH OF CHILDREN
See Side Panel for Additional Cautions

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

NET CONTENTS

Manufactured by

ECONOMY PRODUCTS COMPANY, INC.
SHENANDOAN, IOWA 51601

DIRECTIONS FOR LIVESTOCK PEST CONTROL (Continued)

face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gallons of spray.

Do not use in poultry houses.

For Control of Chorioptic Mites on Dairy and Beef Cattle. Mix 2½ pints of CI-PONA Insecticide Emulsifiable per 16 gallons of water (0.25% concentration). Spray animals thoroughly. Retreat within 10-14 days for lasting control.

WARNING

WARNING! HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

During commercial or prolonged exposure in spray-mixing and loading operations, wear clean rubber gloves and a mask or respirator of a type found suitable by the U. S. Bureau of Mines for protection against CIODRIN-VAPONA insecticide. Use this material only for recommended purposes and at recommended dosages.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not treat animals within one day of slaughter.

Do not apply dermally in combination with Triclorfon.

DECONTAMINATE AND DESTROY USED CONTAINERS. Do not re-use for any purpose. Decontaminate containers by rinsing thoroughly with water and alkaline detergent. Destroy containers by breaking and burying fragments in isolated area. Dispose of rinsings in a way as not to constitute a hazard, or contaminate water supplies.

NOTICE: Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 8770-802
EPA Est. 8770-1A-1
8770-NB-1

8919B-MBEPKO

