




# VMI LIVESTOCK, POULTRY & PREMISE SPRAY

## RABON® 50% WETTABLE POWDER INSECTICIDE

**ACTIVE INGREDIENT:**  
2-chloro-1-(2,4,5-trichlorophenyl)vinyl dimethyl phosphate ..... 50.0%  
**INERT INGREDIENTS** ..... 50.0%  
**Total** ..... 100.0%

BY WEIGHT



### GENERAL DIRECTIONS

RABON® W. P. is suitable for use in conventional power or low pressure knapsack sprayers. Occasional agitation is recommended to prevent undue settling of the suspension. Follow the USE DIRECTIONS for the proper percent solution needed for a specific insect and areas or types of wall surfaces. Refer to DILUTION TABLE for the quantity of insecticide needed to make the percent solution recommended.

### LIVESTOCK USE DIRECTIONS

Animal	Insect	% Solution	Remarks
Beef Cattle	Horn Flies, Lice	0.35%	Apply as coarse spray. Use between 1/2 and 1 gallon of spray per animal depending on size and hair coat. There is no withholding period from last application to slaughter.
	Lone Star Ticks	0.5%	
Swine	Lice	0.5%	Apply as a coarse spray using 1 to 2 quarts per head to thoroughly wet the animal. Repeat in 2 weeks if necessary.

### POULTRY USE DIRECTIONS

Insect	Type of Housing	% Solution	Remarks
Lice & Mites	Wire Cages	0.5%	Apply directly to birds (1 gal./100 birds), spray vent and fluff areas from below. Repeat when necessary. Do not repeat more often than every 14 days. For individual bird treatment apply 1 ounce of the mixture per bird.

**NOTE:** For maximum lasting control of the northern fowl mite, penetration of the feathers around the vent area is absolutely essential. Use power sprayer at 100-125 psi at NO LESS THAN RECOMMENDED RATE. More attention must be given each individual bird when using low-pressure equipment. Treat roosters carefully and thoroughly to avoid reinfestation in breeding flocks.

Floor Management Litter	0.5%	Apply 1-2 Gal./1,000 sq. ft. evenly for penetration to litter surfaces. Also apply thoroughly to walls, roosts, cracks, crevices and interiors. Spray birds lightly.	
Floor Management Litter	50% WP	Treat evenly and thoroughly using 2.5 oz./100 sq. ft.	
Floor Management Dust Box	50% WP	Mix evenly throughout top layer of box contents using 2.5 oz./50 birds.	
Floor Management Roost Paint	1.0%	Treat with brush or spray thoroughly, particularly cracks and crevices using 1 pt./100 ft.	
Fowl Tick (Blue Bug)	All Types	1.0%	Apply 1 gal./100-150 sq. ft. thoroughly to walls, ceiling, floor cracks and crevices with a power sprayer.

TAKE TIME

OBSERVE LABEL DIRECTIONS

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Harmful if swallowed. Avoid breathing spray mist and dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

See Other Cautions on Side Panel

NET CONTENTS \_\_\_\_\_

Packed by  
**ECONOMY PRODUCTS CO., INC.**  
SHENANDOAH, IOWA 51601

Insect	Problem Area	% Solution	Remarks
Flies	Dairy barns, poultry houses, swine barns — other animal buildings.	2.0%	For dry whitewashed wood or concrete block surfaces 1 gal. of solution/500 sq. ft.
		1.0%	For unpainted wood or painted concrete block surfaces 1 gal. of solution/500 sq. ft.
		1.0%	For masonite or galvanized sheet metal surfaces 1/2 gal. of solution/500 sq. ft.
Maggots	Poultry droppings, manure piles, garbage piles, under feed troughs.	1.0%	Apply 1 gal. of mixed solution 100 sq. ft. Penetrate problem area first time—repeat every 7-10 days thereafter.
Ticks	Campgrounds, backyards, picnic areas, recreational parks, other outdoor living areas.	0.5%	Spray infested areas thoroughly (approximately 25 gals. of mixed solution/acre). Spray along foot paths and roadsides leading to such areas.

### DILUTION TABLE

Amount of Rabon® Wettable Powder	AMOUNT OF WATER FOR			
	0.35% Solution	0.5% Solution	1.0% Solution	2.0% Solution
1/2 lb. ....	9 Gal.	6 Gal.	3 Gal.	1 1/2 Gal.
2 lb. ....	36 Gal.	25 Gal.	12 1/2 Gal.	6 Gal.
4 lb. ....	75 Gal.	50 Gal.	25 Gal.	12 1/2 Gal.
8 lb. ....	150 Gal.	100 Gal.	50 Gal.	25 Gal.

### PRECAUTIONS IN USING

**ENVIRONMENTAL CAUTIONS:** This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**DISPOSAL PRECAUTIONS:** Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

**NOTICE:** Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 3770-311  
EPA Est. 3770-NB-1

3653B-PBEPCCO