

ROSE and FLORAL DUST

ACTIVE INGREDIENTS:

Folpet [N-(trichloromethylthio) phthalimide]* 7.50%

Malathion (O,O-Dimethyl dithiophosphate of
diethyl mercaptosuccinate) 4.50%

**A coordination product of zinc ion and manganese
ethylene bisdithiocarbamate 3.25%

In which the ingredients are:

Manganese ++65000

Zinc ++08125

Ethylene bisdithiocarbamate ion
(C₄H₆N₂S₄) = 2.51875

INERT INGREDIENTS: 84.75%

*Phaltan

**U.S. Pat. Nos. 2,504,404; 2,974,156 and Patent Pending

CAUTION
KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional Cautions

HANDY READY TO USE DUST GUN

TO OPERATE —

- (1) Twist tube to break label at perforation.
- (2) Remove tape which covers hole at top of gun.
- (3) Use pumping stroke to spray dust.

ROSE and FLORAL DUST is effective for —

ROSES

To control black spot, powdery mildew, aphid, mites, Japanese beetle (adult), tarnished plant bug, leafhopper, mealybug, thrips.

FLOWERS

Carnations: Rust, alternaria leafspot, aphid, mealybug, thrips.

Chrysanthemums: Septoria leafspot, aphid, four lined leaf bug, tarnished plant bug, leafhopper, thrips.

Aster, Phlox, Snapdragon: powdery mildew, aphid, tarnished plant bug, leafhopper, mealybug, mites.

Zinnias, Marigold: Alternaria leafspot, aphid, Japanese beetle (adult), tarnished plant bug, leafhopper, mealybug, mites.

Snapdragons: Anthracnose, rust, tarnished plant bug, mites.

Iris: Didymellina leafspot, aphid, thrips.

Gladia'us: Botrytis blight, curvularia, aphid, tarnished plant bug, leafhopper, thrips.

Note: Do not use on Boston, Pteris and Maidenhair fern, crassula and canaerti juniper as injury may occur.

DIRECTIONS FOR USE

Application — Dust up and under the leaves thoroughly. Best results will be obtained by dusting in the early morning or evening when the air is calm and temperature is low.

Authorities rightly maintain that the secret of successful disease and pest control depends largely upon the proper timing of the applications. Foliage growth is rapid in the early spring, so that there is a daily increase in leaf surface exposed to insect and fungus attack. Keep this new growth covered by frequent and timely applications. For disease and insect control begin dusting when new growth appears and repeat at 7 to 10 day intervals throughout the growing season. For disease, treat twice weekly during periods of rapid new growth and severe infection caused by continuous rains or high humidity. Such a program is the best insurance against disease and insect damage during the period of primary infection and infestation.

CAUTION: Harmful if inhaled or swallowed. May cause irritation of nose, throat and skin. Avoid inhalation of dust. Avoid contact with skin. Wash thoroughly after handling. Do not contaminate food, water, dishes or utensils, or feedstuffs.

ENVIRONMENTAL CAUTIONS: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of

RECEIVED

ACTIVE INGREDIENTS:

Folpet [N-(trichloromethylthio) phthalimide]* 7.50%

Malathion (O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate) 4.50%

**A coordination product of zinc ion and manganese ethylene bisdithiocarbamate 3.25%

In which the ingredients are:

Manganese ++65000

Zinc ++08125

Ethylene bisdithiocarbamate ion (C₄H₆N₂S₄) =2.51875

INERT INGREDIENTS:84.75%

*Phaltan

**U.S. Pat. Nos. 2,504,404; 2,974,156 and Patent Pending

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional Cautions

NET CONTENTS 3 OZ. NOV 7 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under

Manufactured by

EPA Reg. No.

3770-313

ECONOMY PRODUCTS COMPANY, INC.

SHENANDOAH, IOWA 51601

ROSES

To control black spot, powdery mildew, aphid, mites, Japanese beetle (adult), tarnished plant bug, leafhopper, mealybug, thrips.

FLOWERS

Carnations: Rust, alternaria leafspot, aphid, mealybug, thrips.

Chrysanthemums: Septoria leafspot, aphid, four lined leaf bug, tarnished plant bug, leafhopper, thrips.

Aster, Phlox, Snapdragon: powdery mildew, aphid, tarnished plant bug, leafhopper, mealybug, mites.

Zinnias, Marigold: Alternaria leafspot, aphid, Japanese beetle (adult), tarnished plant bug, leafhopper, mealybug, mites.

Snapdragons: Anthracnose, rust, tarnished plant bug, mites.

Iris: Didymellina leafspot, aphid, thrips.

Gladiolus: Botrytis blight, curvularia, aphid, tarnished plant bug, leafhopper, thrips.

Note: Do not use on Boston, Pteris and Maidenhair fern, crassula and canaerti juniper as injury may occur.

DIRECTIONS FOR USE

Application — Dust up and under the leaves thoroughly. Best results will be obtained by dusting in the early morning or evening when the air is calm and temperature is low.

Authorities rightly maintain that the secret of successful disease and pest control depends largely upon the proper timing of the applications. Foliage growth is rapid in the early spring, so that there is a daily increase in leaf surface exposed to insect and fungus attack. Keep this new growth covered by frequent and timely applications. For disease and insect control begin dusting when new growth appears and repeat at 7 to 10 day intervals throughout the growing season. For disease, treat twice weekly during periods of rapid new growth and severe infection caused by continuous rains or high humidity. Such a program is the best insurance against disease and insect damage during the period of primary infection and infestation.

CAUTION: Harmful if inhaled or swallowed. May cause irritation of nose, throat and skin. Avoid inhalation of dust. Avoid contact with skin. Wash thoroughly after handling. Do not contaminate food, water, dishes or utensils, or feedstuffs.

ENVIRONMENTAL CAUTIONS: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

IMPORTANT: Keep away from fire and sparks. Store in a cool, dry place. Keep container closed when not in use.

DISPOSAL PRECAUTIONS: Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 3770-313
EPA Est. 3770-NB-1

4839B-MBEP CO