

DIRECTIONS

ROACHES, ANTS, SPIDERS, SILVERFISH, CRICKETS, CLOVER MITES, EARWIGS in and about **homes, dairies and food processing plants**, use 2% malathion in water. To make a water solution, dilute 2 tablespoons of 57% malathion emulsifiable in 1 quart of water (1 1/2 ounces per gallon). Apply by means of coarse spray or paint brush to window sills, baseboards, under sinks and stoves, and to other areas frequented by insects. Spray surfaces until wet. Care should be taken to thoroughly treat all cracks and crevices. Water-based sprays may stain finished surfaces. **DO NOT USE IN MILK PROCESSING ROOMS.** Do not use in commercial food preparation areas or in edible products areas of food processing plants. Avoid contamination of food, utensils, working surfaces, containers, and equipment.

POULTRY (chickens, ducks, geese, turkeys): Direct application to poultry for **northern fowl mites** and **poultry lice** mix 2 tablespoons per 1 gallon of water per 100 to 150 birds. Apply sprays for control of **northern fowl mite, poultry lice** and as a supplement to premise treatments of **chicken red mite**. Repeat in 4 to 8 weeks or when necessary.

POULTRY HOUSES: For **northern fowl mite, chicken red mite, poultry lice, flies**, mix 1 tablespoon per 1 gallon of water. Apply spray liberally to **litter, walls, ceilings, roosts, nests and adjacent areas**. Force spray into all cracks and crevices. Repeat as necessary. For poultry ticks mix 6 to 7 oz. per 1 gallon of water. Apply spray liberally to walls, ceilings and adjacent areas. Force spray into cracks and crevices. Repeat as necessary.


When applying sprays avoid contamination of feed, food containers and watering troughs.

ORNAMENTALS, FLOWERS, SHRUBS, SHADE TREES: For **mealybugs, aphids, spider mite, rose leafhopper**, mix 1 1/2 pints per 100 gallons of water and make full cover applications to the point of run-off. For smaller areas, mix 1 1/2 teaspoons per gallon of water. Repeat as necessary. To control bagworms, tent caterpillar, azalea scale, spider mites, leaf scale, soft brown scale, and scurfy scale, mix 2 teaspoonfuls per 1 gallon of water. Do not apply to certain ferns such as Boston, Maidenhair and Pteris, and some Crassula. This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

USDA Reg. No. 3772-7

J-6960A-K

MANUFACTURED FOR
EARL MAY SEED & NURSERY CO.
SHENANDOAH, IOWA



57%

MALATHION SPRAY

KILLS

FLIES, MOSQUITOES and GARDEN INSECTS

ACTIVE INGREDIENTS:

Malathion*	57.0%
Xylene Type Aromatic Hydrocarbon Solvent	32.0%
INERT INGREDIENTS:	11.0%
Total	100.0%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION

KEEP OUT OF THE REACH OF CHILDREN.

See other warnings on side panel.

NET CONTENTS 1 GALLON

DII

FLY AND MOSQUITO CONTROL emulsifiable to 3 gallons rate of 1 gallon per 1,000 sq gal. per 1,000 square feet on or congregate; walls, ceiling. Use a large droplet spray the run-off. For treating floors in thion 57% in 2 gallons of water can be applied to window and doorways, with a sprink on animals. Remove animals, processed.

POTATOES, TOMATOES: For or more applications, as ne malathion 57% emulsifiable ough coverage. For mixing 1 gallon of water. Make no ap harvest. **PEAS, BEANS, PE CUCUMBER, SQUASH, MEL Mites, Leafhoppers, Mexican** per acre in sufficient water. NOT APPLY WITHIN 3 DA cucurbits unless plants are

CAUTION: Harmful if swall Avoid contact with skin; was eating or smoking. Avoid co Keep out of reach of childre may cause cholinesterase in antidotal.

This product is highly toxic or residues on crops. Prote from your Cooperative Agric

Do not reuse empty containe burning. Stay away from smo

NOTICE: Seller warrants tha cal description and is reason label when used in accordanc ditions of use, but neither th of MERCHANTABILITY OR I POSE, express or implied, ex trary to label instructions, or conditions not reasonably for the risk of any such use.

MANUF
EARL MAY SE
SHENA

E-2-72
3772-7



57% MALATHION SPRAY

KILLS
FLIES, MOSQUITOES and GARDEN INSECTS

ACTIVE INGREDIENTS:

Malathion*	57.0%
Xylene Type Aromatic Hydrocarbon Solvent	32.0%
INERT INGREDIENTS:	11.0%
Total	100.0%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION

KEEP OUT OF THE REACH OF CHILDREN.

See other warnings on side panel.

NET CONTENTS 1 GALLON

DIRECTIONS

FLY AND MOSQUITO CONTROL AID: Add 8 oz. of 57% malathion emulsifiable to 3 gallons of water. Apply the spray at the rate of 1 gallon per 1,000 square feet on painted surfaces and 2 gal. per 1,000 square feet on unpainted surfaces where flies alight or congregate: walls, ceiling, stanchions, window sills and fences. Use a large droplet spray that wets all surfaces to the point of run-off. For treating floors in dairy barns, use 5 ounces of malathion 57% in 2 gallons of water. As a spot treatment this mixture can be applied to window frames, stanchions, support beams, and doorways, with a sprinkling can or paint brush. Do not use on animals. Remove animals. Do not use where milk is stored or processed.

POTATOES, TOMATOES: For **aphids** and **leafhoppers**, make one or more applications, as needed, of a spray containing 1 pint malathion 57% emulsifiable per acre in sufficient water for thorough coverage. For mixing small quantities use 1 teaspoonful per gallon of water. Make no application later than 3 days prior to harvest. **PEAS, BEANS, PEPPERS, RUTABAGAS, EGG PLANT, CUCUMBER, SQUASH, MELON, ONION:** For **Aphids, Spider Mites, Leafhoppers, Mexican Bean Beetle, Onion Thrips**, use 1 pt. per acre in sufficient water to insure thorough coverage. **DO NOT APPLY WITHIN 3 DAYS OF HARVEST.** Do not apply to cucurbits unless plants are dry.

CAUTION: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin; wash thoroughly after using, and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep out of reach of children. **NOTE TO PHYSICIAN:** Malathion may cause cholinesterase inhibition. Atropine and 2-PAM are antidotal.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

MANUFACTURED FOR
EARL MAY SEED & NURSERY CO.
SHENANDOAH, IOWA

1, CRICKETS, CLOVER
ies and food processing
make a water solution,
ulsifiable in 1 quart of
means of coarse spray
under sinks and stoves,
pray surfaces until wet,
all cracks and crevices,
faces. DO NOT USE IN
e in commercial food
reas of food processing
asils, working surfaces,

ys): Direct application
ultry lice, mix 2 table-
50 birds. Apply sprays
lice and as a supple-
mite. Repeat in 4 to 8

nite, chicken red mite.
gallon of water. Apply
sts, nests and adjacent
vices. Repeat as neces-
per 1 gallon of water.
d adjacent areas. Force
necessary.

on of feed, food con-

DE TREES: For mealy-
mix 1 1/2 pints per 100
ications to the point of
ons per gallon of water.
tent caterpillar, azalea
scale, and scurfy scale.
Do not apply to certain
ris, and some Crassula.
of any body of water.
his label.

J-6960A-K

DR
RSERY CO.

A

6-2-72
3772-7