

GENERAL DIRECTIONS

Apply 5% SEVIN Dust when insects or their damage appear. Make application in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat at 7 to 10-day intervals or as necessary unless shorter interval is specified below. Do not apply in excess.

VEGETABLE INSECT CONTROL

May be used up to day of harvest on the following: **BEANS** (green beans, lima beans, snap beans, cowpeas, black-eyed): Mexican bean beetle, leafhopper, cucumber beetle, Japanese beetle. **Corn:** corn rootworm adults, European corn borer, fall armyworm, corn earworm. For corn borers, apply thoroughly into the whorls when plant is about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 2 to 3-day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required, depending on severity of infestation. Timing and good coverage are essential for effective control. **CARROTS:** flea beetle, leafhopper, harlequin bug. **CUCUMBER, SUMMER and WINTER SQUASH, MELONS, PUMPKINS:** pickleworm, melonworm, cucumber beetle, squash bug, flea beetle, leafhopper. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. **OKRA:** corn earworm, stink bug. **POTATO** (apply as foliage application only). **TOMATO, EGG-PLANT, PEPPER:** Colorado potato beetle, flea beetle, leafhopper, European corn borer, fall armyworm, lace bugs, tomato fruit worm, tomato hornworm, tarnished plant bugs, stink bugs.

Do not apply within 3 days of harvest on the following: **GARDEN BEETS, TURNIPS, SALSIFY** (Do not apply within 14 days of harvest if tops are to be used as food or feed), **BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI:** flea beetle, harlequin bug, armyworm, cabbage caterpillar, corn earworm. **PARSNIPS, RADISHES, RUTABAGAS:** flea beetle, leafhopper.

Do not apply within 14 days of harvest on the following: **LETTUCE (LEAF), SPINACH, SWISS CHARD, CHINESE CABBAGE, COLLARDS, ENDIVE (ESCAROLE), KALE, MUSTARD GREENS, PARSLEY:** armyworm, flea beetle, leafhopper, cabbage caterpillar.

GRASSHOPPERS ON THE ABOVE CROPS

Apply as often as necessary to control grasshoppers on the above crops.


SMALL FRUIT INSECT CONTROL

Use sufficient dust to obtain full coverage. Apply when insects or their damage appear. Repeat at 7 to 10-day intervals or as necessary.

GRAPES: grape leaf folder, grape leafhopper, and grape leaf skeletonizer. Apply just before first brood leaf folder larvae emerge from rolls and as needed for leafhoppers. May be applied up to day of harvest.

STRAWBERRIES: meadow spittlebug, strawberry leaf roller and strawberry weevil. May be used up to 1 day before picking.

BLUEBERRIES: blueberry maggot, cherry and cranberry fruitworms and Japanese



SEVIN* DUST INSECTICIDE

*For the Control of Certain Insects on
Vegetables, Small Fruit, and Pets*

3-29-71
3722-12

CUCUMBER SUMMER

CARROTS: WINTER SQUASH MELONS PUMPKINS:

POTATO
PLANT PEPPER

OKRA:
TOMATO EGG

Do not apply within 3 days of harvest on the following: GARDEN BEETS, TURNIPS, SALSIFY

BROCCOLI BRUSSELS SPROUTS CABBAGE CAULIFLOWER KOHLRABI PAR SNIPS RADISHES RUTABAGAS:

Do not apply within 14 days of harvest on the following: LETTUCE (LEAF) SPINACH SWISS CHARD CHINESE CABBAGE COLLARDS ENDIVE (ESCAROLE) KALE MUSTARD GREENS PARSLEY:

GRASSHOPPERS ON THE ABOVE CROPS

SMALL FRUIT INSECT CONTROL

GRAPES:

STRAWBERRIES:

BLUEBERRIES:

BLACKBERRIES BOYSENBERRIES LOGANBERRIES AND RASPBERRIES:

DOGS CATS

PET USE

in kennel dog and cat quarters

CAUTION

NOTICE

MANUFACTURED FOR
E. H. MAY SEED & NURSERY CO.

SEVIN DUST INSECTICIDE

*For the Control of Certain Insects on
Vegetables, Small Fruit, and Pets*

ACTIVE INGREDIENTS:	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	85%
INERT INGREDIENTS:	15%
Total	100%

*SEVIN is a registered trade mark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

CAUTION

KEEP OUT OF THE REACH OF CHILDREN.

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Do not Take Internally. Avoid Breathing of Dust. Skin Contact May Be Harmful. Avoid Skin Contact. Wash Hands Before Eating. Avoid Contamination Of Food and Feedstuffs. Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

SEE OTHER CAUTIONS ON BACK PANEL